# CAUTION

**KEEP OUT OF REACH OF CHILDREN**
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**TITAN**

# DFF 25 + MCPA 250
## SELECTIVE HERBICIDE

**ACTIVE CONSTITUENTS: 250g/L MCPA present as the ETHYL HEXYL ESTER**
**25g/L DIFLUFENICAN**
**SOLVENTS: 313g/L LIQUID HYDROCARBONS**
**150g/L N-METHYL-2-PYRROLIDONE**

GROUP 12 | 4 HERBICIDE



For the control of certain broadleaf weeds in winter cereals and clover as specified in the Directions For Use table.

APVMA Approval No.: 65650/150571V

**Pack Size: 5L; 20L; 110L; 1000L**


TITAN AG
CROP PROTECTION

TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | titanag.com.au


**EC** Formulation Type **Emulsifiable Concentrate**

**IN A TRANSPORT EMERGENCY**
**• DIAL 000 • POLICE OR FIRE BRIGADE**

9 3082

**ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (contains diflufenican and MCPA, 2-ethylhexyl ester)**
**PG III Hazchem •3Z**

**DIRECTIONS FOR USE**

**Restraints:** DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.
DO NOT apply to crops under stress due to disease or insect damage.
DO NOT apply to frost-affected crops or if frosts are imminent.
DO NOT apply when heavy rain is expected within 4 hours.

**SPRAY DRIFT RESTRAINTS**

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.
DO NOT allow bystanders to come into contact with the spray cloud.
DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.
DO NOT apply unless the wind speed is between 3 and 20 km/h at the application site during the time of application.
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

| CROP | WEEDS CONTROLLED | STAGE OF WEED GROWTH | STATE | RATE/ha | CRITICAL COMMENTS |
|---|---|---|---|---|---|
| **Cereals** Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)<br><br>**Pasture** Newly sown and established clover based pasture, clover for hay and seed production | Wild Radish | Up to the 2 leaf stage and not more than 60mm in diameter | WA only | 250mL | **CROP STAGE**<br>**Cereals**<br>Up to 750mL (3 leaf to fully tillered stage - Z13 to 30).<br>Over 750mL (5 leaf to fully tillered stage - Z15 to 30). Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).<br>**WA only:** DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).<br>**Warning:** TITAN DFF 25 + MCPA 250 Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions).<br>**Clover**<br>Application is recommended prior to the eighth trifoliate leaf stage, however, applications prior to the third leaf stage may result in crop damage especially under stressed conditions and in sandy soils. DO NOT apply to Annual Medics or Lucerne.<br>**Warning:** TITAN DFF 25 + MCPA 250 Selective Herbicide may cause transient crop yellowing of clover, and may affect growth and seed set of some varieties of Clover. (Refer to "Crop Tolerance" section of General Instructions).<br>**WEED STAGE**<br>Apply when weeds are actively growing. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as:<br>* high crop and weed density;<br>* late season germinations;<br>* abnormal weed growth (including early flowering), higher rates of product (up to the maximum rate of application specified for that weed) may be required.<br>TITAN DFF 25 + MCPA 250 Selective Herbicide will not effectively control:<br>* regrowth of suppressed weeds;<br>* transplanted weeds;<br>* regrowth from rhizomes or roots;<br>* weeds growing under stress from previous herbicide applications.<br>**GRAZING**<br>Efficacy on larger weeds will be improved by grazing with normal levels of stock after the 7 day withholding period. Refer to 'Protection of Livestock' for grazing precautions.<br>**APPLICATION**<br>Activity of this product will be reduced if weeds are stressed. Optimum results will be obtained if good soil moisture exists at and after application. Where crop or weed density is high, water volume should be increased.<br>**WILD RADISH**<br>TITAN DFF 25 + MCPA 250 Selective Herbicide will provide residual control of Wild Radish for up to 4 weeks after application. Effective residual activity of this product may be reduced where:<br>* rates lower than 1.0L/ha are used;<br>* dry conditions prevail;<br>* poor coverage of the soil surface is achieved;<br>* crop is planted in non-wetting sand;<br>* soils contain a high content of organic matter.<br>Optimum results will be obtained if good soil moisture exists at and after application. |
| | | Up to the 4 leaf stage and not more than 120mm in diameter | ALL STATES | 500mL | |
| | | Up to the 6 leaf stage and not more than 150mm in diameter | | 750mL | |
| | | Up to the 8 leaf stage and not more than 180mm in diameter | | 1.0L | |
| | Charlock, Hedge Mustard, Indian Hedge Mustard, Shepherd's Purse, Turnip Weed, Wild Turnip | Up to the 2 leaf stage and not more than 60mm in diameter | | 500mL | |
| | | Up to the 4 leaf stage and not more than 120mm in diameter | | 750mL | |
| | | Up to the 6 leaf stage and not more than 150mm in diameter | | 1.0L | |
| | London Rocket | Up to the 5 leaf stage and not more than 120mm in diameter | QLD only | 750mL | |
| | Ward's Weed | | SA only | | |
| | Capeweed | Up to the 2 leaf stage and not more than 60mm in diameter | ALL STATES | 500mL | |
| | | Up to the 4 leaf stage and not more than 120mm in diameter | | 1.0L | |
| | Crassula | Up to the 2 leaf stage | | 500mL | |
| | | Up to the 4 leaf stage | | 750mL | |
| | Prickly Lettuce | Up to the 2 leaf stage | | 500mL | |
| | | Up to the 4 leaf stage | | 750mL | |
| | | Up to the 6 leaf stage | | 1.0L | |
| | Dense-flower Fumitory | Up to the 2 leaf stage | | 750mL | |
| | Corn Gromwell, Saffron Thistle, Toad Rush | | | 1.0L | |
| | Deadnettle | | NSW, VIC, SA only | | |
| | Sorrel | Up to the 2 leaf stage | VIC only | 1.0L | |
| | Canola (rape) | Up to the 4 leaf stage | ALL STATES | 500mL | |
| | Purple Goosefoot | Up to the 6 leaf stage | QLD only | 500mL | |
| | Turnip Weed, Wild Turnip | Cotyledon to 2 leaf stage | NSW only (West of Newell Hwy) SA only (Eyre peninsula north of the line between Venus Bay and Cowell) | 350mL | |



TITAN AG CROP PROTECTION

| CROP | WEEDS CONTROLLED | STAGE OF WEED GROWTH | STATE | RATE/ha | CRITICAL COMMENTS |
|---|---|---|---|---|---|
| **Cereals** Wheat, Barley, Oats, Triticale, Cereal Rye | Fumitory | 2-6 leaf stage | ALL STATES | 500mL + 200mL Terbutryn (500g/L) | **Refer also to all Critical Comments relating to weed stage, grazing, application and Wild Radish above.**<br>* Reduced efficacy (suppression only) may be achieved on Wild Radish larger than 8 leaf or greater than 180mm in diameter.<br>DO NOT use this tank mix if Cereals are undersown with Lucerne or Annual Medics.<br>**Crop Stage**<br>TITAN DFF 25 + MCPA 250 Selective Herbicide 350mL + MCPA LVE 200mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30).<br>TITAN DFF 25 + MCPA 250 Selective Herbicide 500mL + MCPA LVE 200mL: Apply from 3 leaf to fully tillered (Zadok's Z13 to Z30).<br>TITAN DFF 25 + MCPA 250 Selective Herbicide 500mL + MCPA LVE 400mL: Apply from 5 leaf stage to fully tillered (Zadok's Z15 to Z30).<br>Optimum results are achieved when sprayed at 3-5 leaf crop stage (generally 4-8 weeks post sowing).<br>**WA only:** DO NOT apply to Barley or Kulin Wheat before the 5 leaf stage (Z15).<br>**Warning:** DFF 25 + MCPA 250 Selective Herbicide may cause transient crop yellowing of cereals. Some varieties of oats have not been tested. (Refer to "Crop Tolerance" section of General Instructions).<br>Observe instructions also on MCPA LVE product label. |
| **Cereals** Wheat, Barley, Oats, Triticale, Cereal Rye (including cereals undersown with clover)<br><br>**Pasture** Newly sown and established clover based pasture, clover for hay and seed production | **SUPPRESSION OF THE FOLLOWING WEEDS** | | | | |
| | Saffron Thistle | Up to the 6 leaf stage | ALL STATES | 1.0L | |
| | Chickweed, Fireweed, Hexham Scent (King Island Melilot), Iceplant, Mouse-eared Chickweed, Nightscented Stock, Paterson's Curse, Peppercress, Skeleton Weed, Long Storksbill, Volunteer Lupins | Up to the 4 leaf stage | | | |
| | Wireweed (Hogweed) | | | 750mL | |
| | Common Sowthistle (Milk Thistle), Cowvine, Dock, Doublegee (Spiny Emex), Fat Hen, Horehound, Hyssop Loosestrife, Marshmallow, Rough Poppy, Scarlet Pimpernel, Stemless Thistle, Tree Hogweed, Variegated Thistle, Vetch (Tares) | Up to the 2 leaf stage | | 1.0L | |
| **Cereals** Wheat, Barley, Oats, Triticale, Cereal Rye | Wild Radish | Up to the 4 leaf stage and not more than 120mm in diameter | ALL STATES | 350mL plus 200mL MCPA LVE (500g/L) | |
| | | Up to the 6 leaf stage and not more than 150mm in diameter | | 500mL plus 200mL MCPA LVE (500g/L) | |
| | | Up to the 8 leaf stage and not more than 180mm in diameter* | | 500mL plus 400mL MCPA LVE (500g/L) | |

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS**
**Crop Harvest: NOT REQUIRED WHEN USED AS DIRECTED.**
**All Crops: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**



TITAN AG CROP PROTECTION

**WEED LIST**

| Common name | Scientific name |
|---|---|
| Canola (rape) | *Brassica napus* |
| Capeweed | *Arctotheca calendula* |
| Charlock | *Sinapis arvensis* |
| Chickweed | *Stellaria media* |
| Common Sowthistle (Milk Thistle) | *Sonchus oleraceus* |
| Corn Gromwell | *Buglossoides arvense* |
| Cowvine | *Ipomoea lonchophylla* |
| Crassula | *Crassula* spp. |
| Deadnettle | *Lamium amplexicaule* |
| Dense-flower Fumitory | *Fumaria densiflora* |
| Dock | *Rumex* spp. |
| Doublegee (Spiny Emex) | *Emex australis* |
| Fat Hen | *Chenopodium album* |
| Fireweed | *Senecio* spp. |
| Fumitory | *Fumaria* spp. |
| Hedge Mustard | *Sisymbrium officinale* |
| Hexham Scent (King Island Melilot) | *Melilotus indicus* |
| Horehound | *Marrubium vulgare* |
| Hyssop Loosestrife | *Lythrum hyssopifolia* |
| Iceplant | *Mesembryanthemum* spp. |
| Indian Hedge Mustard | *Sisymbrium orientale* |
| London Rocket | *Sisymbrium irio* |
| Long Storksbill | *Erodium botrys* |
| Marshmallow | *Malva parviflora* |
| Mouse-eared Chickweed | *Cerastium glomeratum* |
| Night-scented Stock | *Matthiola longipetala* |
| Paterson's Curse | *Echium plantagineum* |
| Peppercress | *Lepidium* spp. |
| Prickly Lettuce | *Lactuca serriola* |
| Purple Goosefoot | *Scleroblitum atriplicinum* |
| Rough Poppy | *Papaver hybridum* |
| Saffron Thistle | *Carthamus lanatus* |
| Scarlet Pimpernel | *Anagallis arvensis* |
| Shepherd's Purse | *Capsella bursa-pastoris* |
| Skeleton Weed | *Chondrilla juncea* |
| Sorrel | *Rumex* spp. |
| Stemless Thistle | *Onopordum acaulon* |
| Toad Rush | *Juncus bufonius* |
| Tree Hogweed | *Polygonum patulum* |
| Turnip Weed | *Rapistrum rugosum* |
| Variegated Thistle | *Silybum marianum* |
| Vetch (Tares) | *Vicia sativa* |
| Volunteer Lupins | *Lupinus* spp. |
| Ward's Weed | *Carrichtera annua* |
| Wild Radish | *Raphanus raphanistrum* |
| Wild Turnip | *Brassica tournefortii* |
| Wireweed (Hogweed) | *Polygonum aviculare* |

**GENERAL INSTRUCTIONS**

**TOLERANCE**

Some pre-emergence herbicides, such as atrazine, can cause stress to certain crops resulting in an increase in crop damage when using this product. Subterranean Clover is particularly sensitive.

**CEREALS**

After application some transient crop yellowing may occur. This usually appears as yellow or white banding on leaves. Provided the crop is not under stress from pre-emergent herbicide, root disease, insect damage, frost, dry or excessively moist conditions, the development of the crop and subsequent growth will be unaffected.

**Warning (Oats):** The tolerance of oat varieties Esk and Nile (the 2 main varieties grown in Tasmania) to TITAN DFF 25 + MCPA 250 Selective Herbicide has not been tested. Test a small area of crop before using TITAN DFF 25 + MCPA 250 Selective Herbicide over large areas. Consult your local TITAN AG representative for advice on specific varieties.

**PASTURE**

The tolerance of clover varieties to TITAN DFF 25 + MCPA 250 Selective Herbicide can vary with rate of application, soil type, crop health, stage of growth and degree of moisture and temperature stress.

**Warning:** TITAN DFF 25 + MCPA 250 Selective Herbicide may result in transient crop yellowing and suppression of growth with a resultant initial reduction in dry matter, particularly at rates in excess of 500mL/ha and in areas of double spray. For this reason we recommend application prior to the 8 trifoliate leaf stage. However, at the lower rates (500mL/ha and less) and under normal growing conditions, subsequent growth and seed yield should not be affected. Under normal growing conditions, the following varieties have shown acceptable levels of foliage tolerance to TITAN DFF 25 + MCPA 250 Selective Herbicide applied at 500mL/ha:

**ArrowLeaf:** Zulu

**Balansa:** Paradana

**Berseem:** Sacromonte

**Persian:** Kyambro, Lupers, Maral

**White:** Haifa

**Subterranean Clover:** Daliak, Dalkeith, Denmark, Esperance, Geraldton, Goulburn, Karridale, Larissa, Leura, Mt Barker, Nungarin, Rosedale, Seaton Park, Trikkala and Woogenellup.

Some pasture grasses, including Phalaris and Cocksfoot, may show some initial reduction in vegetative growth after application of TITAN DFF 25 + MCPA 250 Selective Herbicide. Care should be exercised if sensitive clover varieties or grasses are included in the pasture sward.

Varieties not listed should be tested before using TITAN DFF 25 + MCPA 250 Selective Herbicide over large areas. Consult your local TITAN AG representative for advice on specific varieties.

**SUBSEQUENT CROPS**

To reduce effect on subsequent susceptible crops (eg. Canola), ensure thorough cultivation of soil prior to the sowing of these crops.

**MIXING**

To ensure even mixing, half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly then add the remainder of the water. Agitate again before spraying commences. Reseal part-used product container immediately after use. Spray mixtures containing TITAN DFF 25 + MCPA 250 Selective Herbicide should not be left to stand overnight. Prolonged periods of exposure to cold temperatures could result in settling out of the product in the mixture.

**Warning:** The rubber components present in some spraying units may be affected by exposure to the solvents in TITAN DFF 25 + MCPA 250 Selective Herbicide and some other agricultural products. To reduce this risk it is recommended that the spray unit be thoroughly washed with a boom cleaner and fresh water after use. Contact the spray unit manufacturer to determine the suitability of the rubber components for use with agricultural products.

**APPLICATION**

**Boom Sprayer:** A minimum of 50L of water per hectare should be used, however, for optimum results water rates of 70-100L/ha are recommended. Increase the water volume if weed infestation is heavy or crop cover is dense. Complete coverage of weeds is essential.

**Aircraft (NSW, VIC, SA only):** Apply in a minimum of 30L water per hectare. Effective weed control will only be achieved where good coverage of leaf surface is achieved.



TITAN AG CROP PROTECTION

**COMPATIBILITY**
The following products are physically compatible with TITAN DFF 25 + MCPA 250 Selective Herbicide as a two-way mixture in the spray tank but should only be used for the crops specified:

| Crop | TITAN DFF 25 + MCPA 250 Selective Herbicide | Compatible Product |
|---|---|---|
| Wheat, Triticale and Cereal Rye only | Up to 750mL/ha | TITAN Diclofop-Methyl 500 EC Selective Herbicide (also Barley) |
| Cereals (including undersown) | All rates | TITAN Chlorpyrifos 500 Insecticide, TITAN Dimethoate 400 Insecticide |
| Cereals (not undersown) | Up to 500mL/ha | TITAN Metsulfuron 600 WG Herbicide, TITAN Chlorsulfuron 750 WG Herbicide, TITAN MCPA LVE, TITAN Triasulfuron 750 WG Herbicide |
| | All rates | TITAN Bromoxynil 200 Selective Herbicide, TITAN Amine 625 Selective Herbicide, TITAN Clopyralid 300 Herbicide, TITAN Dicamba 700 WG Herbicide (up to 115g only) |
| Wheat, Barley, Triticale, and Cereal Rye only (not undersown) | | TITAN Tralkoxydim 400 WG Herbicide |
| Wheat only (not undersown) | | TITAN Clodinafop 240 EC Herbicide |
| Clover | Up to 750mL/ha | TITAN Quizalofop-P-Ethyl 200 Herbicide, TITAN Fluazifop 212 Post-emergence Selective Herbicide |
| Subterranean Clover | | TITAN Simazine 500 Flowable Herbicide, TITAN Simazine 500 + TITAN Paraquat 250 mixture |
| | Up to 1.0L/ha | TITAN 2,4-DB 500 SL Herbicide |

When mixing with other herbicides, crop yellowing may be enhanced.
When mixing with TITAN Diclofop-Methyl 500 EC Selective Herbicide, Wildcat or Tristar Advance some reduction in the efficacy and speed of action of these products may occur. If the crop is stressed, the application of the herbicide tank-mixtures may cause yield reduction.
When mixing with TITAN Dicamba 700 WG Herbicide a temporary wilting may be evident in some crops after application.
Growers should seek advice before spraying recently released cereal varieties. Use the recommended rates for both herbicides in the tank mixture as well as the surfactant recommendation of the grass herbicide. If another herbicide is applied as a tank mix, observe the plantback restrictions on that label.
DO NOT add surfactant when mixing TITAN DFF 25 + MCPA 250 Selective Herbicide and TITAN Metsulfuron 600 WG Herbicide.

**Simazine:** Refer to the simazine label for correct application rates, especially with regard to soil types. This product may be mixed in the spray tank with one of the following insecticides according to the directions for use on this product: TITAN Glufosinate 200 Herbicide, TITAN Alpha-Cypermethrin 250 SC Insecticide, TITAN Lambda-Cyhalothrin 250 CS Insecticide and TITAN Bifenthrin 100 Insecticide/Miticide.

**Warning:** DO NOT use crop oils with Titan DFF 25 + MCPA 250 Selective Herbicide or TITAN DFF 25 + MCPA 250 Selective Herbicide tank mixtures with other products in Cereals. As formulations of other manufacturer's products are beyond the control of TITAN AG Pty Ltd, all mixtures should be tested prior to mixing commercial quantities.

**RESISTANT WEEDS WARNING**
GROUP 12 4 HERBICIDE
TITAN DFF 25 + MCPA 250 Selective Herbicide is a member of the phenoxy and nicotinanilide groups of herbicides and acts by inhibiting carotenoid biosynthesis at the phytoene desaturase step (PDS inhibitors) and disrupting plant cell growth. For weed resistance management TITAN DFF 25 + MCPA 250 Selective Herbicide is both a Group 12 and Group 4 herbicide. Some naturally occurring weed biotypes resistant to TITAN DFF 25 + MCPA 250 Selective Herbicide and other Group 12 and Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by TITAN DFF 25 + MCPA 250 Selective Herbicide or other Group 12 and Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN DFF 25 + MCPA 250 Selective Herbicide to control resistant weeds.

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**
DO NOT apply under weather conditions, or from spraying equipment, that may cause drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

**PROTECTION OF LIVESTOCK**
**Grazing Precaution:** Sprayed weeds may become more palatable to stock and a higher intake of some weeds may result in stock poisoning and death from causes such as nitrate poisoning.
Care should be taken especially where Capeweed, Paterson's Curse and Variegated Thistles predominate in the pasture. Avoid grazing with young or breeding stock.
DO NOT graze horses or pigs on Paterson's Curse. If in doubt, contact your nearest Department of Agriculture.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**
Dangerous to fish. DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

**STORAGE AND DISPOSAL**
**5L, 20L:** Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the empty containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**Envirodrum Micro Matic Valve (110L):** Store the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal.
DO NOT contaminate the Envirodrum with water or any foreign matter. After each use of the product, please ensure that the Micro Matic coupler delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the Envirodrum have been used, please return the Envirodrum to the point of purchase. The Envirodrum remains the property of TITAN AG Pty Ltd.

**1000L:** Store in the closed original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use then the integrity of this product cannot be assured. If this occurs TITAN AG Pty Ltd should be notified immediately. This minibulk/bulk container is reusable and remains the property of TITAN AG Pty Ltd. DO NOT rinse empty container. Empty contents fully into application equipment. Close all valves and return to the point of supply for refill or storage. No other liquid, solid or pesticide product should be put into it. When empty return to TITAN AG Pty Ltd for cleaning, relabelling and refilling.

**SAFETY DIRECTIONS**
Harmful if swallowed. Will damage the eyes. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

**FIRST AID**
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.

**SAFETY DATA SHEET**
Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN DFF 25 + MCPA 250 Selective Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au



TITAN AG CROP PROTECTION

**CONDITIONS OF SALE:** TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Combustible liquid. Harmful if swallowed. Causes skin irritation. Causes serious eye damage. May cause respiratory irritation. Suspected of causing cancer. May damage the unborn child. Very toxic to aquatic life with long lasting effects. Precautionary Statements: Do not handle until all safety precautions have been read and understood. Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Use only outdoors or in a well-ventilated area. Wear protective gloves, protective clothing, eye protection and face protection. Avoid breathing mist/vapours/ spray. Wash all exposed external body areas thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. IF exposed or concerned: Get medical advice/ attention. Immediately call a POISON CENTER/doctor/physician/first aider. In case of fire: Use alcohol resistant foam or normal protein foam to extinguish. Collect spillage. IF SWALLOWED: Call a POISON CENTER/doctor/physician/first aider if you feel unwell. IF ON SKIN: Wash with plenty of water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Rinse mouth. If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Store locked up. Store in a well-ventilated place. Keep container tightly closed. Dispose of contents/container to authorised hazardous or special waste collection point in accordance with any local regulation.





TITAN AG
CROP PROTECTION