

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

# CONFIDE 350 SC

## INSECTICIDE

ACTIVE CONSTITUENT: 350g/L IMIDACLOPRID

GROUP **4A** INSECTICIDE

For the control of various insect pests of cotton, fruit, vegetables, ornamentals and turf as a foliar spray and as a soil-applied treatment for the control of various canegrubs in sugarcane, certain pests in apples and citrus, and silverleaf whitefly in various vegetable crops and an injection treatment for control of certain pests in bananas as specified in the Directions for Use table.

APVMA Approval No.: 86992/137278

Pack Size: 1L; 5L; 10L; 20L; 110L; 200L; 1000L



TITAN AG Pty Ltd | ABN 57 122 081 574  
15/16 Princes Street, Newport NSW 2106  
Tel (02) 9999 6655 | Fax (02) 9999 0483  
titanag.com.au

IN A TRANSPORT EMERGENCY

# DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING  
NOT A DANGEROUS GOOD  
ACCORDING TO THE AUSTRALIAN  
DANGEROUS GOODS (ADG) CODE  
FOR TRANSPORT BY ROAD AND RAIL

## DIRECTIONS FOR USE

### SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at [apvma.gov.au/spraydrift](http://apvma.gov.au/spraydrift)

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 15 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

### Restraints:

DO NOT use on crops produced hydroponically or in glasshouses and other covered situations.

DO NOT apply TITAN Confide 350 SC Insecticide or any other Group 4A Insecticide as a foliar spray after soil application of TITAN Confide 350 SC Insecticide in that crop.

DO NOT apply more than one soil application of TITAN Confide 350 SC Insecticide or any other soil applied Group 4A Insecticide per crop for Vegetables or Sugarcane, per season for Citrus or per two years for Apples.

DO NOT apply more than one application of TITAN Confide 350 SC Insecticide or any other Group 4A Insecticide per crop for Bananas.

1. FOLIAR APPLICATIONS			
CROP	PEST	RATE	CRITICAL COMMENTS
Brassicas	Grey Cabbage Aphid, Turnip Aphid	15mL/100L or 170mL/ha	Apply at first sign of aphid infestation. Add a wetting agent.
Capsicum	Green Peach Aphid		Apply at first sign of aphid infestation.
Cotton	Cotton Aphid	145mL/ha + Pulse Penetrant at 0.2% v/v (2mL/L water) or equivalent organosilicone surfactant (check with TITAN AG for recommended organosilicone surfactants)	<p>The addition of Pulse Penetrant or equivalent is critical for the performance of TITAN Confide 350 SC Insecticide. Apply early in the establishment of an aphid infestation when numbers are low (ie no more than 1 or 2 leaves per plant with honeydew present). Applications made later than this may result in reduced control.</p> <p>Shorter residual control may be evident and a repeat application of a registered aphicide (follow the Cotton Insecticide Resistance Management Strategy for Cotton Aphid) may be required to achieve complete control:</p> <ul style="list-style-type: none"> <li>• If applications of TITAN Confide 350 SC Insecticide plus Pulse are timed too late (see above); or</li> <li>• If existing high density aphid colonies (hotspots) are present; or</li> <li>• If aphids have established throughout the plant canopy (especially lower in the canopy); or</li> <li>• If there is high reinfestation pressure; or</li> <li>• If there is rapid crop growth; or</li> <li>• If TITAN Confide 350 SC Insecticide plus Pulse is used following a spray-failure (eg. resistance to organophosphate or carbamate insecticides).</li> </ul> <p><b>Note:</b> Where resistance to carbamates or organophosphates is suspected, TITAN Confide 350 SC Insecticide plus Pulse should be used first so as not to delay control of the aphids present.</p> <p>Aphids treated with TITAN Confide 350 SC Insecticide plus Pulse may still be present on the plant but will not be feeding. Control of aphids should initially be assessed by a reduction in fresh honeydew and not on the presence of aphids on the plant. After ingesting TITAN Confide 350 SC Insecticide, aphids may take up to 5 days to die.</p>
	Mirids, Brown Flea Beetle		Apply when pest numbers reach treatment threshold levels as determined by field checks.
Cucumber	Silverleaf Whitefly, including type B	15mL/100L or 145mL/ha	Apply at first sign of whitefly infestation. Apply dilute sprays (15mL/100L) to run-off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Cucurbits	Green Peach Aphid	15mL/100L or 170mL/ha	Apply at first sign of aphid infestation.
Duboisia	Green Peach Aphid	15mL/100L	Apply when aphid numbers reach spray threshold levels as determined by regular monitoring. Ensure thorough coverage of all leaves.
Eggplant	Green Peach Aphid	15mL/100L or 170mL/ha	Apply at first sign of aphid infestation.
	Melon Thrips	15mL/100L or 145mL/ha	Apply at first sign of Melon Thrips infestation. Apply dilute sprays (15mL/100L) to run-off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Ornamental Plants	Aphids, Azalea Lace Bug, Bronze Orange Bug, Harlequin Bug, Citrus Mealybug, Greenhouse Thrips, Fullers Rose Weevil	15mL/100L	Apply as a thorough cover spray at first sign of insect infestation.
	Hibiscus Flower Beetle	30mL/100L	Spray buds and flowers as needed.
	Longtailed Mealybug	30mL/100L + surfactant	Apply 3 sprays 2 weeks apart. Use a non-ionic surfactant at label rate.
	Psyllids	15mL/100L	Spray at first sign and then a week later.
	Soft Scales		Spray in late spring or when small scales are first seen. Apply 3 sprays 2 weeks apart. Add a wetting agent.

CROP	PEST	RATE	CRITICAL COMMENTS
Pandanus Trees	Flatid ( <i>Jamella australiae</i> )	<b>Spot Spray</b> 500mL/100L of water <b>Stem Injection</b> 1L/1L of water	<b>Spot Spray:</b> Spray 100mL of mixture directly into the leafy throat of each head. <b>Stem Injection:</b> Drill holes 0.5-1cm in diameter and 10cm deep at an angle of 30° and 1-1.5m above ground level. Drill one hole per limb (or trunk in single trunked trees). Apply 5mL of mixture in each hole and seal the hole. DO NOT reapply in the same holes. Uptake of TITAN Confide 350 SC Insecticide and therefore control of the pest in heavily infested heads already showing severe damage, will be slow and may be incomplete.
Potato	Green Peach Aphid	15mL/100L or 170mL/ha	Apply at first sign of aphid infestation.
Roses	Aphids	15mL/100L	Apply as a thorough cover spray at first sign of aphid infestation.
Stone Fruit	Green Peach Aphid, Black Peach Aphid	<b>Dilute spraying</b> 15mL/100L <b>Concentrate spraying</b> Refer to the Mixing/ Application section	Apply at first sign of aphid infestation. Apply as a full cover spray, ensuring thorough coverage. Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use in equipment that requires rates greater than 75mL/100L of water (ie greater than 5 X concentrate).
Sweet Potato	Silverleaf Whitefly, including type B	15mL/100L or 145mL/ha	Apply at first sign of whitefly infestation. Apply dilute sprays (15mL/100L) to run-off. Ensure thorough coverage of underside of leaves. Use of droppers will improve coverage of underside of leaves.
Tomato	Green Peach Aphid	15mL/100L or 170mL/ha	Apply at first sign of aphid infestation.
Turf	First instar larvae of: Black Beetle, Argentinian Scarab, Pruinose Scarab	1.5L/ha or 15mL/100m <sup>2</sup> Spray with at least 400L water per hectare to ensure even coverage. Preferably spray onto wet or dewy grass. Irrigate with 12mm of water commencing within 1 hour of application.	Apply at peak egg hatch, which is mid-spring to mid-summer depending on species.
	Larvae of Billbug		Monitor adult activity through late spring and early summer. Spray when numbers peak or when small larvae (4mm) are found in the thatch or surface soil. Early application is essential to minimize grass damage due to feeding.
2. SOIL/OTHER METHODS OF APPLICATION			
CROP	PEST	RATE	CRITICAL COMMENTS
Apples	Woolly Aphid	<b>Chemical control</b> 3.5mL/1L of water/tree	For trees up to 7 years of age. During late summer or autumn, Apple trees with Woolly Aphid colonies or damage should be identified and marked for treatment the following season. At green tip to petal fall, apply 1 litre of the prepared TITAN Confide 350 SC Insecticide mixture to moist soil immediately around the base of the tree trunk. Ensure the mixture infiltrates the soil around the trunk and does not run-off the soil. Control weeds before application. DO NOT disturb or remove the soil around the trunk during the season. If aerial colonies are present at application, maximum effectiveness may not be achieved until the following season. DO NOT treat more than once in any 2 year period.
		<b>Beneficial insect plus chemical control</b> (eg. <i>Aphelinus mali</i> plus TITAN Confide 350 SC Insecticide) 1.75mL/1L of water/tree	
Azaleas in pots	Azalea Lace Bug	2mL/250mL water/pot	Use as a soil drench for pots up to 20L capacity. Prior to application remove mulch and dead vegetation, and moisten the soil surface. Apply the TITAN Confide 350 SC Insecticide mixture and then water it in well immediately after application.
Bananas – Cavendish	Banana Rust Thrips (Nth QLD, NT, Nth WA only)	2.5 or 3.5mL/stool Inject undiluted, or diluted 50:50 with water	The higher rate may give improved control in some situations. <b>Application Timing:</b> DO NOT inject bunched plants. Select the best follower and inject into the base. DO NOT inject any follower more than once. Injection can occur at any time within 3 months after harvest of the mother plant or nurse-suckering. To limit the risk of plant damage from injection, inject only those followers that are at least 1.5m tall to the throat of the plant. Plants smaller than 1.5m tall to the throat can be treated but there is significant risk of plant injury from injection. For application in plant bananas the same minimum height requirement applies as above, plus the plant should be at a stage at least 3 months prior to bell emergence. <b>Application Method:</b> Injection should occur 15cm from the base of the plant at an off-centre, downward angle to a depth of 5 to 10cm. DO NOT inject into the centre of the plant as this may result in plant death. Injection should be conducted in a manner which ensures applied chemical is retained within the pseudostem. Irrigation or moderate to heavy or prolonged periods of rainfall may saturate the internal structure of the pseudostem, leading to 'run-out' from injection holes. Allow sufficient time following irrigation and rainfall events, before application, to minimise such 'run-out'. TITAN Confide 350 SC Insecticide should be used as part of an integrated pest management approach which should include the use of other measures for control of Banana Rust Thrips, such as bell injection and bunch applications. <b>Note:</b> Application during conditions conducive to Banana Spider Mite may enhance population development. Under these conditions, continue to monitor mite populations following TITAN Confide 350 SC Insecticide application, taking appropriate action where thresholds are exceeded.
	Banana Weevil Borer (QLD, NSW, NT, WA only)	2.5mL/stool Inject undiluted, or diluted 50:50 with water	
Bananas – Lady Finger (QLD, NSW, NT, WA only)	Banana Weevil Borer	2.5 or 3.5mL/stool Inject undiluted, or diluted 50:50 with water	

CROP	PEST	RATE	CRITICAL COMMENTS
Brassicacae (Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kohlrabi)	Silverleaf Whitefly, including type B	<b>Plant hole drench, furrow spray</b> 35mL/1000 seedlings	<p><b>Plant hole drench, furrow spray</b>  <b>Green Peach Aphid, Onion Thrips:</b> When TITAN Confide 350 SC Insecticide is used for the control of Silverleaf Whitefly, including type B, control of Green Peach Aphid and Onion Thrips will also be achieved. TITAN Confide 350 SC Insecticide provides effective management of pest populations. However, TITAN Confide 350 SC Insecticide may not provide complete control of pests for the entire growing period. Crops should be monitored for pests following planting and throughout the life of the crop. If pests are observed in the crop additional chemical control may be required, in which case an insecticide with a different mode of action should be used. Refer to GENERAL INSTRUCTIONS for resistance management strategy information.</p> <p><b>Plant hole drench (at or post-transplanting):</b> Mix TITAN Confide 350 SC Insecticide with sufficient water to allow a constant volume of at least 50mL of drench mixture per plant. Apply the selected volume of drench mixture in the planting hole at planting or within 2 days after planting. At planting and when applying drench, steps should be taken to ensure workers do not contact treated soil or drench mixture.</p> <p><b>Furrow spray (prior to planting):</b> Mix TITAN Confide 350 SC Insecticide with water, using at least 2 litres of spray mixture per 100m of row. Apply to open furrow not earlier than 5 days prior to planting as a narrow band of spray centred under the plant row. DO NOT leave the sprayed band exposed to sunlight; sprayed soil should be covered immediately. After final shaping of the planting bed, the treated layer of soil should be approximately 100mm below the soil surface. This method of application may provide less effective control than seedling drench or plant hole drench application methods. At planting, steps should be taken to ensure workers do not contact treated soil.</p>
Capsicums, Cucurbits, Eggplant, Sweet Potato, Tomatoes	Silverleaf Whitefly, including type B	14mL/100 metres of row	<p><b>Sub-surface trickle irrigation injection:</b> Apply once only 5-7 days after planting (or 5-7 days from seed emergence if planted from seed). Begin injection only after water has reached the furthest drip points and soil is partially wetted up. After TITAN Confide 350 SC Insecticide injection is completed, continue irrigation only until lines are flushed, not longer than 1 hour.</p> <p>DO NOT apply TITAN Confide 350 SC Insecticide using surface trickle irrigation or any other type of above ground irrigation system. Subsequent irrigations should occur only when soil moisture measurements indicate the need for addition of water. DO NOT over irrigate or cause run-off. In situations where root development in the crop is slow, evidence of control may be delayed.</p>
		14mL/100 metres of row (Mix with water, using at least 2 litres of spray mixture per 100m row)	<p><b>Furrow spray pre-plant:</b> Apply to open furrow not earlier than 5 days prior to planting as a narrow band of spray centred under the plant row.</p> <p>DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. Sprayed soil should be covered immediately. After final shaping of the planting bed, the treated layer of soil should be approximately 100mm below the soil surface. At planting, steps should be taken to ensure workers do not contact treated soil.</p>
Capsicums, Eggplant, Tomatoes		14mL/100 metres of row (Mix with sufficient water, to allow a constant volume of at least 50mL of drench mixture per plant)	<p><b>Plant hole drench:</b> DO NOT apply TITAN Confide 350 SC Insecticide by this method where plant spacing along the row exceeds 60cm. Apply the selected volume of drench mixture in the planting hole at planting or within 2 days after planting. Steps should be taken to ensure workers do not contact treated soil or drench mixture.</p>
Citrus	Black Citrus Aphid, Citrus Leafminer, Pink Wax Scale, Red Scale	9mL/tree Apply as a soil drench* OR via micro-sprinkler* or drip irrigation*	<p>* Refer to Application directions under GENERAL INSTRUCTIONS for detailed information on application methods. Application via micro-sprinkler and drip application methods will give best results when used in conjunction with other control methods such as the release of beneficial species (eg. parasitic wasps). Treatment is only recommended for trees of up to 4m in height.</p> <p><b>Citrus Leafminer, Black Citrus Aphid:</b> Apply TITAN Confide 350 SC Insecticide in late spring after main flowering has finished (October to December) prior to the summer or autumn flush. Apply prior to pest establishment or at the first signs of pest infestation.</p> <p><b>Red Scale, Pink Wax Scale:</b> Monitor crop throughout late spring to early summer (October to December). If scale is observed, apply TITAN Confide 350 SC Insecticide after main flowering has finished and prior to or at the onset of crawler emergence.</p> <p><b>Multiple flowering and/or overlapping cropping:</b> Where extended flowering and/or multiple flowering periods occur eg. Lemons and Limes, or if the previous seasons crop is still hanging on the tree during or at the end of a new seasons flowering (overlapping cropping) eg. Valencia Oranges, TITAN Confide 350 SC Insecticide should only be applied:</p> <ul style="list-style-type: none"> <li>• When there is a minimum of 20 weeks to the next harvest and,</li> <li>• According to the timing for specific pests described above and,</li> <li>• After the previous crop has been harvested or stripped and,</li> <li>• When the main flowering period has finished.</li> </ul> <p>DO NOT apply more than once per season.</p>

CROP	PEST	RATE	CRITICAL COMMENTS
Citrus (non-bearing only)	Citrus Leafminer	6mL/tree Apply as a soil drench* OR via drip irrigation*	<p>* Refer to Application directions under GENERAL INSTRUCTIONS for detailed information on application. For application by drip, the emitters should be in close proximity to the base of the tree to maximise contact with the tree root system and to minimise the opportunity for breakdown of TITAN Confide 350 SC Insecticide by UV exposure, otherwise TITAN Confide 350 SC Insecticide uptake and hence Leafminer control may be reduced.</p> <p>Treatment is only recommended for vegetative (non-bearing) trees. Apply TITAN Confide 350 SC Insecticide between late spring to late summer (October to February) prior to a Leafminer susceptible growth flush. Apply prior to pest establishment or at the first signs of pest infestation. If longer residual control is desired then follow the recommendation above for the use of the product on Citrus at the 9mL/tree rate. DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. DO NOT apply more than once per season.</p>
Elm	Elm Leaf Beetle	4mL/25mm of tree diameter at breast height	Mix the required dose in sufficient water to adequately treat each tree. Use at least 50L of mix per tree up to a tree diameter of 400-500mm and then 100L per tree for larger trees. Inject mix to a depth of 20-30cm in a minimum of 4 injection sites per tree, 0.75 to 1.5m apart, arranged in an evenly spaced grid to just beyond the drip line. Ensure root zone is adequately moist with active root growth. Keep treated area moist for 7-10 days after treatment. Treat at least 6-10 weeks prior to pest attack in late winter or early spring when roots are active. DO NOT treat if soil is waterlogged.
Ornamentals in pots	Scarab Beetle Larvae	2mL/5L water	Use as a soil drench. 5L of mixture will treat twenty 6L pots. Prior to application remove mulch and dead vegetation and moisten the soil surface. Apply the TITAN Confide 350 SC Insecticide mixture and then water it in well immediately after application.
Peanuts (QLD, NSW, NT only)	Greyback Canegrub, Childers Canegrub, Negatoria Canegrub, French's Canegrub, Southern One-Year Canegrub, Peanut White Grub	6.5-13mL per 100 metres of Peanut row (equivalent to 0.71L-1.43L/ha for single Peanut row with 0.91m spacing between rows)	Apply at planting as a narrow in-furrow band spray directed on or below the Peanut seed. DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. Ensure that product is immediately covered by soil. Canegrub and White Grub pest thresholds have not been established in Peanuts, however, the use of the highest rate where heavy grub infestations are expected eg. above 1 grub per plant, is recommended. Refer to Application directions under GENERAL INSTRUCTIONS.
Potatoes	Silverleaf Whitefly, including type B	14mL/100 metres of row (Mix with water using 1.5- 3.0 litres of spray mixture per 100m of row)	<b>Furrow spray at planting:</b> Apply as part of the planting process to the open furrow as a narrow spray band (100-150mm wide) centred in the plant row at seed level. If seed piece breakdown is considered to be a risk, eg. in hot sandy soils, avoid wetting the seed during application of TITAN Confide 350 SC Insecticide. DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. Sprayed soil should be covered immediately. After the planting operation, the treated soil should be at least 100mm below the surface.
	Green Peach Aphid	9mL/100 metres of row (Mix with water using 1.5- 3.0 litres of spray mixture per 100m of row)	
Roses	Aphids	2mL/2L water/plant	Use as a soil drench by pouring mixture evenly around drip zone. Use this rate for plants up to 1m high. For each additional metre of plant height, add 1.2mL extra of TITAN Confide 350 SC Insecticide to the 2L of water. Prior to application remove mulch and dead vegetation and moisten the soil surface. Apply the TITAN Confide 350 SC Insecticide mixture and then water it in well immediately after application.
Sugarcane (Ratoon cane only)	Greyback Canegrub	16-22mL per 100 metres of cane row (equivalent to 1.05-1.44L/ha for single row cane with 1.52m spacing between rows)	<p>Apply from September to November to fields which are at high risk of Greyback Grub damage. In areas where early flights of beetles occur, application should be early within this period. Late applications where large 3rd instar larvae dominate the grub population will not be as effective. Application should be made while stools are small enough to avoid excessive damage. Use the high rate when high grub populations are expected; eg. an average greater than 4 grubs per stool. Apply only as a subsurface soil application behind coulters (refer to Application directions under GENERAL INSTRUCTIONS). Soil should have moisture at coulter depth at the time of application or should receive at least 15mm of rainfall or irrigation within 1 week.</p> <p>DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. After application ensure the TITAN Confide 350 SC Insecticide treated band is covered by at least 100mm of soil and that coulter slits are filled in completely. DO NOT apply more than once per season.</p>
	Childers Canegrub	11-16mL per 100 metres of cane row (equivalent to 0.72-1.05L/ha for single row cane with 1.52m spacing between rows)	<p>Apply from September to November. Check for the presence of grubs from September onwards. Apply TITAN Confide 350 SC Insecticide when grub numbers reach an economic threshold (about 3 grubs per stool). Use the high rate when grub populations are high, eg. an average greater than 5 grubs per stool or if application is late (damage already visible). Apply only as a subsurface soil application behind coulters (refer to Application directions under GENERAL INSTRUCTIONS). Soil should have moisture at coulter depth at the time of application or should receive at least 15mm of rainfall or irrigation within 1 week.</p> <p>DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. After application ensure the TITAN Confide 350 SC Insecticide treated band is covered by at least 100mm of soil and that coulter slits are filled in completely. DO NOT apply more than once per season.</p>



CROP	PEST	RATE	CRITICAL COMMENTS
Sugarcane (Ratoon cane) All Sugarcane areas	French's Canegrub	16-22mL per 100 metres of cane row (equivalent to 1.05-1.44L/ha for single row cane with 1.52m spacing between rows)	Begin monitoring for the presence of grubs in September, and continue to monitor at regular intervals. Apply TITAN Confide 350 SC Insecticide immediately if grub numbers have reached an economic threshold (about 3 grubs per stool). Early applications are more effective than later ones. DO NOT apply later than November. Use the high rate when grub populations are high, eg. an average greater than 5 grubs per stool, or if application is late (damage already visible). Apply only as a subsurface soil application behind coulters. Refer to Application directions under GENERAL INSTRUCTIONS.
	Negatoria Canegrub	11-16mL per 100 metres of cane row (equivalent to 0.72-1.05L/ha for single cane row with 1.52m spacing between rows)	
Sugarcane (Ratoon cane) Southern Queensland (Bundaberg and South) and NSW only	Childers Canegrub, Plectris Canegrub, Rhopaea Canegrub, Southern One-Year Canegrub	11-16mL per 100 metres of cane row (equivalent to 0.72-1.05L/ha for single cane row with 1.52m spacing between rows)	Begin monitoring for the presence of grubs in September (December for Southern One-Year Canegrub) and continue to monitor at regular intervals. Apply TITAN Confide 350 SC Insecticide if grub numbers reach an economic threshold (eg. 3-4 grubs per stool). Early applications are more effective than later ones. Use the high rate when grub populations are high eg. an average greater than 5 grubs per stool or if application is late (damage already visible). Apply only as a subsurface soil application behind coulters. Refer to Application directions under GENERAL INSTRUCTIONS.
Sugarcane (Plant cane) All Sugarcane areas	Greyback Canegrub	16-22mL per 100 metres of cane row (equivalent to 1.05-1.44L/ha for single row cane with 1.52m spacing between rows)	<b>Moderate – high pressure (2 or more grubs/stool expected). August – November applications from planting to hilling-up.</b> Apply at planting, or at first working, or at half-open drill stage, or at fill-in or final hilling. Apply at final hilling only if sufficient soil cover can be applied. Use the high rate where heavy Canegrub infestation is expected. Apply in a narrow spray band in the planted row. Treated soil must be covered immediately. Refer to Application directions under GENERAL INSTRUCTIONS.
		11mL per 100 metres of cane row (equivalent to 720mL/ha for single row cane with 1.52m spacing between rows)	<b>Low pest pressure (less than 2 grubs/stool expected). September – November applications at fill-in and hilling-up only.</b> Apply at fill-in or final hilling only. Apply at final hilling only if sufficient soil cover can be applied. Apply in a narrow spray band in the planted row. Treated soil must be covered immediately. Refer to Application directions under GENERAL INSTRUCTIONS.
Sugarcane (Plant cane) Southern Queensland (Bundaberg and South) and NSW only	Childers Canegrub, Negatoria Canegrub, Plectris Canegrub, Rhopaea Canegrub, Southern One-Year Canegrub	11-16mL per 100 metres of cane row (equivalent to 0.72-1.05L/ha for single row cane with 1.52m spacing between rows)	Apply generally in spring or summer either at planting, or at first working, or at half-open drill stage, or at fill-in or final hilling. Apply at final hilling only if sufficient soil cover can be applied. If larvae are likely to be present at or soon after planting (eg. in a plough out/replant situation) then early application is recommended. If larvae pressure is not anticipated until the crop is established, then application at the later recommended timing ie closer to the first Canegrub larvae presence may be more effective; for example, autumn plantings, application should generally be delayed until spring. Use the high rate where heavy Canegrub infestation is expected, or to obtain longer residual activity. Apply in a narrow spray band in the planted row. Treated soil must be covered immediately. Refer to Application directions under GENERAL INSTRUCTIONS.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

#### WITHHOLDING PERIODS

##### HARVEST

**Brassicas, Capsicum, Eggplant, Potatoes, Sweet Potatoes: DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION**

**Cotton: DO NOT HARVEST FOR 13 WEEKS AFTER APPLICATION**

**Cucurbits (foliar): DO NOT HARVEST FOR 1 DAY AFTER APPLICATION**

**Tomatoes (foliar): DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION**

**Stone Fruit: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION**

**Citrus: DO NOT HARVEST FOR 20 WEEKS AFTER APPLICATION**

**Sugarcane: DO NOT HARVEST FOR 21 WEEKS AFTER APPLICATION**

**All other crops: NOT REQUIRED WHEN USED AS DIRECTED**

##### GRAZING

**Peanuts: DO NOT GRAZE OR CUT TREATED AREAS FOR STOCKFOOD FOR 12 WEEKS AFTER APPLICATION**

**Sugarcane: DO NOT GRAZE OR CUT FOR STOCKFEED FOR 21 WEEKS AFTER APPLICATION**

**All other crops: DO NOT graze any treated area or cut for stock food or feed produce harvested from treated area to animals, including poultry**

##### EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with TITAN Confide 350 SC Insecticide. If you are growing edible produce for export, please check with TITAN AG Pty Ltd for the latest information on MRLs and import tolerances before using TITAN Confide 350 SC Insecticide.

## GENERAL INSTRUCTIONS

### INSECTICIDE RESISTANCE MANAGEMENT STRATEGY (VEGETABLES)

DO NOT apply more than one soil application of TITAN Confide 350 SC Insecticide to each crop. DO NOT use TITAN Confide 350 SC Insecticide or any other Group 4A Insecticide as a foliar spray after soil application of TITAN Confide 350 SC Insecticide in that crop. Refer to district advice for local Silverleaf Whitefly resistance management strategies.

### NOTE FOR VEGETABLE CROPS

Potential migration of Silverleaf Whitefly from neighbouring crops. Adult Silverleaf Whitefly (SLWF) are controlled when they ingest a lethal dose of active ingredient by feeding on a TITAN Confide 350 SC Insecticide treated crop. However, in some very susceptible crops and varieties, the first migration of large populations of adults from adjacent fields may result in significant feeding damage to the crop, although further reproduction and development of the pest will be prevented. To help prevent such damage it is important to minimise the migration of adult SLWF into a treated crop, eg. by applying a 'clean-up' fast-acting insecticide to recently harvested crops. Consideration of factors such as planting sequences and timing, wind direction, variety selection, and general crop hygiene should also be integral to SLWF management. Crops should be monitored for SLWF adult numbers after application of TITAN Confide 350 SC Insecticide, and appropriate fast acting insecticides for control of adults should be applied if economic thresholds are reached or excessive adult feeding damage is observed.

### MIXING

Prior to pouring, shake container vigorously, then add the required amount of TITAN Confide 350 SC Insecticide to water in the tank while stirring or with agitators in motion. TITAN Confide 350 SC Insecticide requires constant agitation in the tank.

### APPLICATION

For all application instructions, refer to the Critical Comments in the Directions for Use Table. Additional instructions for Stone Fruit, Cotton, Sugarcane and Citrus follow.

#### Special Instructions for Stone Fruit

##### Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

##### Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

##### EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500L/ha
2. Your chosen concentrate spray volume: For example 500L/ha
3. The concentration factor in this example is:  $3 \times$  (i.e.  $1500L \div 500L = 3$ )
4. As the dilute label rate is 15mL/100L for stone fruit, then the concentrate rate becomes  $3 \times 15$ , that is 45mL/100L of concentrate spray.

- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- DO NOT use a concentrate rate higher than that specified in the Critical Comments.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**Special Instructions for Cotton:** Thorough coverage of Cotton plants is essential to achieve maximum performance from TITAN Confide 350 SC Insecticide plus Pulse. Equipment should be calibrated to achieve a minimum of 60 droplets/cm<sup>2</sup> on the target foliage. A droplet size for optimum performance from TITAN Confide 350 SC Insecticide plus Pulse is dependent

on equipment and is defined below. DO NOT apply when unfavourable environmental conditions may reduce the quality of spray coverage.

**Ground Application (Cotton):** Application using ground equipment should be made using hollow cone nozzles with a minimum spray volume of 100L/ha. Hollow cone nozzles are recommended but if flat fan nozzles are used, higher water volumes will be required and nozzles should be configured to ensure thorough coverage. A FINE droplet must be used. Where multiple nozzles per row are used, they should be of the same specification to ensure that each nozzle contributes an equal proportion of the required dose. Where multiple nozzles per row are used (particularly for banded applications) ensure the correct nozzle overlap pattern is achieved on the target foliage. Banded applications less than 100% are not recommended beyond the 15 node crop stage.

**Aerial Application (Cotton):** Apply in a minimum spray volume of 25L/ha. A FINE droplet must be used. DO NOT exaggerate swath width or exceed a swath width of 20 to 22m. DO NOT apply TITAN Confide 350 SC Insecticide plus Pulse using Ultra Low Volume (ULV) methods. The use of large droplet placement equipment is not recommended. DO NOT apply treatments by aircraft in situations where drift onto sensitive crops, pastures or desirable trees is likely to occur.

### Application (Sugarcane)

**Plant Cane:** Application can occur at various stages from planting until hilling up as described below. TITAN Confide 350 SC Insecticide should be applied into the plant drill in a narrow band centred on the cane row. Mix TITAN Confide 350 SC Insecticide with water and spray the mixture into the cane drill as a narrow 50-100mm wide band in the middle of the drill. Apply in no less than 1.5L of water per 100 metres of cane row (equivalent to no less than 100L/ha water for single row cane with 1.52m spacing between rows). Adjust the application equipment so that the treated band of soil is at least 50mm above the level of the setts in the soil, and is immediately covered by at least 50mm more soil on top of that layer. There should be at least 100mm of soil cover over the treated layer after the final cultivation or hilling-up operation. The placement of the spray band can be achieved in several ways. Not all application methods are suited to every circumstance. Application at planting is not recommended for control of Greyback Canegrubs in situations where very deep planting is practised.

**At Planting:** Fix a spray nozzle or nozzles on the planter so that the TITAN Confide 350 SC Insecticide spray band is directed at soil just after it covers the setts to a depth of at least 50mm. The boards on the planter need to be adjusted so that another 50mm of soil comes over the top of the treated band of soil. Subsequent cutaway cultivations must not disturb soil to the depth of the treated band.

**At first working, or at half-open drill, or at fill-in:** Apply TITAN Confide 350 SC Insecticide as a spray band 50-100mm wide directed at the base of the young cane shoots so that the middle of the drill is treated, not the sides. A single nozzle per row or a directed nozzle each side of each row may achieve this. The treated band must be covered with at least 50mm of soil immediately. Subsequent cultivations must not disturb soil to the depth of the treated band. There should be at least 100mm of soil cover over the treated layer after the final cultivation or hilling-up operation.

**At hilling up:** Apply TITAN Confide 350 SC Insecticide as twin narrow spray bands 50mm wide on each side of the cane rows directed at the base of the cane rows directed at the base of the young cane shoots. Directed nozzles each side of the row may achieve this. Direct the spray nozzles to ensure that the spray is not blocked by cane shoots or leaves. The treated bands must be covered with at least 100mm of soil immediately as the final step in the shaping of the row profile.

**Ratoon Cane:** Apply only as a subsurface soil application. Apply in no less than 1.5 litres of water per 100 metres of cane row (equivalent to no less than 100L/ha of water for single row cane with 1.52m spacing between rows). Soil should have moisture at coulter depth at the time of application or should receive at least 15mm of rainfall or irrigation within 1 week. DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight. After application, ensure the TITAN Confide 350 SC Insecticide treated band is covered by at least 100mm of soil and that coulter slits are filled in completely.

**Twin coulter method:** Apply subsurface behind twin coulters to a depth of 100-125mm. Coulters should be spaced 220mm to 500mm apart, one either side of the centre of the stool. Coulter slits should be in the sides or top of the stool mound rather than at the base. Space coulters as closely as possible, within the above range, but wide enough to avoid excessive stool damage. Narrow spacings may not be possible with advanced ratoon growth. DO NOT apply using narrow spacings if ratoon growth is advanced such that excessive crop damage from equipment may result.



**Single coultter method ("stool split"):** Apply subsurface behind a single coultter in the centre of the stool. DO NOT apply using this method if ratoon growth is advanced such that excessive crop damage from equipment may result. This method may not be suitable for all areas and has not been fully evaluated in some parts of southern Queensland and NSW. Before using this method, consult your local TITAN AG representative.

**Application (Peanuts):** Fix a spray nozzle or nozzles on the planter so that the TITAN Confide 350 SC Insecticide spray band is directed in a narrow band to soil at the bottom of the planting furrow on or below the peanut seed depth. Apply in no less than 1L of water per 100 metres of peanut row (equivalent to no less than 110L/ha of water for peanuts with 0.91m spacing between rows). There must be at least 50mm soil cover over seed immediately after application. DO NOT leave TITAN Confide 350 SC Insecticide exposed to sunlight.

**Application (Citrus):** Ensure treatment area below canopy is weed free prior to application and remains weed free throughout season, otherwise reduced control will occur.

**Soil drench:** Apply TITAN Confide 350 SC Insecticide in a water mix using 1L of water per tree. Apply evenly to moist soil immediately around the base of the tree trunk to a maximum trunk height of 20cm. Ensure the mixture infiltrates the soil around the trunk and does not run-off the soil. DO NOT disturb or remove the treated soil around the trunk during the season. Irrigation for up to 1 hour (depending on soil type) immediately after a soil drench application is preferred but not essential. DO NOT leave TITAN Confide 350 SC Insecticide mixture exposed to sunlight.

**Drip or micro-sprinklers (below canopy):** For treatment under non-bearing trees, application via micro-sprinklers is not recommended and the Critical Comments in the Directions for Use table should be consulted for further guidance on application by drip/trickle. Drip emitters placed under the tree canopy or near the trunk provide better results than those spaced irregularly or further from the tree. Micro-sprinklers should only be used when the wetting zone can be contained beneath the tree canopy, especially in young orchards where there is minimal canopy and root development. Begin injection only after soil below the trees furthest from the injection point is partially wetted up and irrigation system has reached operating pressure. After TITAN Confide 350 SC Insecticide injection is completed, and lines have been flushed, continue irrigation for no longer than 1 hour. On sandy soils, this irrigation period after lines have been flushed should not be more than 10 minutes. Allow 48 hours before subsequent irrigations. This application technique is best suited to an IPM system, where beneficial species (eg. parasitic wasps) are released. DO NOT disturb or remove the treated soil during the season. DO NOT leave TITAN Confide 350 SC Insecticide mixture exposed to sunlight.

#### PRECAUTION

**Application in Bananas:** Ensure application equipment does not leak. To reduce the risk of exposure from accidental leakage, wear gloves during application.

#### INSECTICIDE RESISTANCE WARNING

For insect resistance management TITAN Confide 350 SC Insecticide is a Group 4A Insecticide. Some naturally occurring insect biotypes resistant to TITAN Confide 350 SC Insecticide and other group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if TITAN Confide 350 SC Insecticide or other Group 4A insecticides are used repeatedly. The effectiveness of TITAN Confide 350 SC Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Confide 350 SC Insecticide to control resistant insects. TITAN Confide 350 SC Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, TITAN AG Pty Ltd representative or local agricultural department agronomist.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers. Confide is toxic to certain aquatic species. Application should be planned to avoid run-off within 48 hours of application. Application should not be made to wet/waterlogged soils. Application is not to be performed if heavy rains are expected to occur within 48 hours. Irrigation run-off from treated areas should be prevented from entering drains and waterways. DO NOT over irrigate or cause run-off. Irrigation should only occur when soil moisture measurements indicate the need for addition of water.

**Citrus:** For the first 48 hours after application irrigation should be restricted to only that which is recommended immediately after application (refer to Application section).

**Sugarcane:** Irrigation should not occur within 48 hours of application.

**Vegetables (furrow spray pre-plant and plant hole drench applications):** Irrigation within 48 hours of application should be minimal, and sufficient to reduce seedling stress only.

**Run-off management:** DO NOT apply within 3 metres of aquatic areas. The growth of a vegetative filter strip between the application site and any water body would also assist.

#### PROTECTION OF LIVESTOCK

Avoid application by surface drippers or micro sprinklers while bees are foraging in the orchard, especially on hot days, as bees can use these devices as sources of water for the hive.

#### STORAGE AND DISPOSAL

Store in the closed, original containers in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycle or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in accordance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if swallowed. Repeated exposure may cause allergic disorders. Wash hands after use. When preparing product for use, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield. After each day's use wash gloves, face shield and contaminated clothing.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131 126; New Zealand 0800 764 766.

#### SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Confide 350 SC Insecticide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au

**CONDITIONS OF SALE:** TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if swallowed. Very toxic to aquatic life with long lasting effects. Precautionary Statements: Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. Rinse mouth. Collect spillage. Dispose of contents/container in accordance with local/regional/national regulations.

