

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

IMMENSE

HERBICIDE

ACTIVE CONSTITUENTS: 33g/L IMAZAMOX present as the AMMONIUM SALT
15g/L IMAZAPYR present as the AMMONIUM SALT

GROUP **2** HERBICIDE

For the early post-emergence control of certain annual grass and broadleaf weeds in imidazolinone-tolerant Wheat, imidazolinone-tolerant Barley, imidazolinone-tolerant Canola, as specified in the Directions For Use section of this label.

APVMA Approval No.: 88704/128638

Pack Size: 1L-1000L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | Fax (02) 9999 0483
titanag.com.au

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL

DIRECTIONS FOR USE

Restrictions:

Apply ONLY to certified imidazolinone-tolerant Wheat, imidazolinone-tolerant Barley and imidazolinone-tolerant Canola varieties with the imidazolinone-tolerance technology, excluding varieties CL STL and CL JNZ. DO NOT apply to conventional or other herbicide tolerant Canola, Wheat and Barley varieties.

DO NOT apply to crops that are stressed due to conditions such as waterlogging, too little moisture, frost, disease or nutritional disorders.

DO NOT apply by aircraft.

DO NOT apply if rain is expected within 2 hours of application. DO NOT apply more than once per season to any one crop.

DO NOT use in imidazolinone-tolerant Wheat crops in tank mix or sequentially with diuron, sulfonamides or sulfonamides.

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the BCPC Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product MUST make an accurate written record of the details of each spray application within 24 hours following application and KEEP this record for a minimum of 2 years. The spray application details that must be recorded are: 1 date with start and finish times of application; 2 location address and paddock/s sprayed; 3 full name of this product; 4 amount of product used per hectare and number of hectares applied to; 5 crop/situation and weed/pest; 6 wind speed and direction during application; 7 air temperature and relative humidity during application; 8 nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application; 9 name and address of person applying this product. (Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic or wetland areas including aquacultural ponds within 50 metres downwind from the application area.

DO NOT apply if there are non-target vegetation or animal habitat within 5 metres downwind from the application area.

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Imidazolinone-tolerant Wheat DO NOT use on CL STL and CL JNZ wheat varieties Imidazolinone-tolerant Barley	Brome (<i>Bromus diandrus</i> and <i>B. rigidus</i>), Barley (<i>Hordeum vulgare</i>), Barley Grass (<i>Hordeum leporinum</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Muskweed (<i>Myagrum perfoliatum</i>), Oats (<i>Avena sativa</i>), Triticale (<i>Triticosecale</i> spp.), Wheat (<i>Triticum aestivum</i>) Non-imidazolinone-tolerant varieties Wild Oats (<i>Avena fatua</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>)	375 to 750mL/ha plus Hasten [†] or Kwickin [†] at 0.5L/100L spray volume	NOT for use on Imidazolinone-tolerant Wheat varieties CL STL and CL JNZ. Apply to Imidazolinone-tolerant Wheat crops from the 3 leaf stage to 1st node stage (Z31). Apply to Imidazolinone-tolerant Barley crops from the 5 leaf stage to 1st node stage (Z31). DO NOT apply to imidazolinone-tolerant Wheat or imidazolinone-tolerant Barley after the 1st node stage (Z31). Applications should be targeted at grass weeds when the majority are in the 2-4 leaf stage and only when within the recommended crop stages. Application to multi-tillered crops may impair weed control because of poor contact and coverage of weeds. – See Compatibility. Tank mixes with 750g/L Clopyralid formulations, eg TITAN Clopyralid 750 SG Herbicide at 60mL/ha will provide control of composite and legume weeds. Tank mixes with 570g/L MCPA formulations, eg TITAN LVE MCPA 570 Herbicide at 500mL/ha will provide control of composite and brassicaceous weeds. The control of brassicaceous weeds will depend on the status of Group 2 resistance in the population. The addition of MCPA formulation will improve control and provide an additional mode of action for resistance management. If other weeds require control, apply appropriate herbicides at least two weeks before or after TITAN Immense Herbicide and only when signs of regrowth or renewed vigour appear, otherwise the effects of the early treatment may affect the performance of the subsequent treatment.
	Annual Ryegrass (<i>Lolium rigidum</i>), Dense Flower Fumitory (<i>Fumaria densiflora</i>), Marshmallow (<i>Malva parviflora</i>), Sub-clover (<i>Trifolium subterraneum</i>) Suppression only: Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver Grasses (<i>Vulpia bromoides</i> and <i>V. myuros</i>)	600 to 750mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	Weed species will either be controlled or suppressed. In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced. The control of Annual Ryegrass varies from excellent to poor depending on the status of Group 2 resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants/m ² , or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of a suitable pre-emergent should be made prior to sowing. A follow up grass selective herbicide may also be necessary.
Canola varieties with imidazolinone-tolerance technology only	Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Muskweed (<i>Myagrum perfoliatum</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>)	300 to 500mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	Read Follow Crop comments and restrictions on the label prior to use. Apply to crop at the 2 to 6 leaf stage. DO NOT apply after the 6 leaf stage. Apply to actively growing weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage. Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage to ensure better contact and coverage. If other weeds require control, apply appropriate herbicides at least two weeks after TITAN Immense Herbicide and only when signs of regrowth or renewed vigour appear, or the effects of TITAN Immense Herbicide may affect their performance.

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Canola varieties with imidazolinone-tolerance technology only – continued	As above plus: Capeweed (<i>Arctotheca calendula</i>), Field Pea (<i>Pisum sativum</i>), Narrow Leaf Lupin (<i>Lupinus angustifolius</i>), Sub-clover (<i>Trifolium subterraneum</i>)	300 to 500mL/ha plus Lontrel [†] Advanced Herbicide at 75 to 150mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	Refer to Critical Comments for 300 to 500mL/ha alone. Lontrel Advanced Herbicide aids in the control of legume and composite weed species. Refer to the Lontrel Advanced Herbicide label. Use rates above 75mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage. Lontrel Advanced Herbicide above 75mL/ha can slightly impair grass control. The addition of Lontrel Advanced Herbicide does not affect the control of Wild Radish and Wild Turnip. (Refer to the Compatibility section of this label and the Lontrel Advanced Herbicide label for further details of use).
	Non-imidazolinone-tolerant Barley (<i>Hordeum vulgare</i>), Barley Grass (<i>Hordeum leporinum</i>), Great Brome (<i>Bromus diandrus</i>), Indian Hedge Mustard (<i>Sisymbrium orientale</i>), Oats (<i>Avena sativa</i>), Rigid Brome (<i>Bromus rigidus</i>), Marshmallow (<i>Malva parviflora</i>), Sub. Clover (<i>Trifolium subterraneum</i>) Non-imidazolinone-tolerant Wheat (<i>Triticum aestivum</i>), Wild Oats (<i>Avena fatua</i>), Wild Radish (<i>Raphanus raphanistrum</i>), Wild Turnip (<i>Brassica tournefortii</i>) <u>Suppression only:</u> Annual Ryegrass ³ (<i>Lolium rigidum</i>), Bedstraw spp. (<i>Galium tricornutum</i> and <i>G. aparine</i>), Doublegee (<i>Emex australis</i>), Silver Grasses (<i>Vulpia bromoides</i> and <i>B. myuros</i>)	600 to 750mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	Apply to crop at the 2 to 6 leaf stage. DO NOT apply after the 6 leaf stage. Apply to actively growing grass weeds in the 3-leaf to 2-tiller stage and broadleaf weeds in the 2 to 6 leaf stage. Use the higher rate when weed numbers are high or towards the upper end of the recommended growth stages, or when the crop is at the 5 to 6 leaf stage, to ensure better contact and coverage. Weeds will either be killed in high numbers (weeds controlled) or in moderate numbers (weeds suppressed). In both cases, surviving plants will be stunted and will be uncompetitive with the crop, and seed set will be prevented or greatly reduced. If other weed species require control, apply appropriate herbicides at least two weeks after TITAN Immense and only when signs of regrowth or renewed vigour appear or the effects of TITAN Immense may affect their performance. ³ The control of Annual Ryegrass varies from excellent to poor depending on the status of Group 2 resistance in the population and environmental conditions. Where the population is expected to exceed 200 plants per sqm, or a high level of control is required, or the ryegrass is known to be resistant or thought to be developing resistance, an application of a pre-emergent herbicide should be made prior to sowing. A tank mix with a grass selective herbicide may also be necessary.
	As above plus Capeweed (<i>Arctotheca calendula</i>), Field Pea (<i>Pisum sativum</i>), Narrow Leaf Lupin (<i>Lupinus angustifolius</i>)	600 to 750mL/ha plus Lontrel Advanced Herbicide at 75 to 150mL/ha plus Hasten or Kwickin at 0.5L/100L spray volume	Refer to Critical Comments for 600 to 750mL/ha alone. DO NOT apply after the 6 leaf stage. Lontrel Advanced Herbicide aids in the control of legume and composite weed species. Refer to the Lontrel Advanced Herbicide label. Use rates above 75mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage. Lontrel Advanced Herbicide above 75mL/ha can slightly impair grass control. The addition of Lontrel Advanced Herbicide does not affect the control of other weeds controlled by TITAN Immense Herbicide. (Refer to the Compatibility section of this label and the Lontrel Advanced Herbicide label for further details of use).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

GRAZING

IMIDAZOLINONE-TOLERANT WHEAT AND IMIDAZOLINONE-TOLERANT BARLEY: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

IMIDAZOLINONE-TOLERANT CANOLA: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.

HARVEST

ALL CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

TITAN Immense Herbicide is for use in imidazolinone-tolerant Wheat, imidazolinone-tolerant Barley and imidazolinone-tolerant Canola. Varieties with imidazolinone-tolerance technology are those that have been bred specifically to be tolerant to TITAN Immense Herbicide.

TITAN Immense Herbicide is absorbed through the leaves, green stems and roots of susceptible weeds and moved from the point of contact throughout the plant. Weeds will either die or will remain stunted and will not compete with the crop. Symptoms of kill may take one to two weeks to develop with death occurring up to one month from treatment. Symptoms first appear at the growth points where young foliage becomes discoloured and distorted before dying. TITAN Immense Herbicide is primarily a post-emergence product. Best results will be achieved when good contact and coverage of weeds occurs and weeds are actively growing. The product must be mixed with Hasten or Kwickin as per the Directions for Use.

TITAN Immense Herbicide also has some residual soil activity under good soil moisture conditions. Residual effects on weeds can be reduced when dry soil conditions follow application before the herbicide has moved to the root zone. Best results will be achieved when application is made to moist soil or if approximately 10mm rain follows within several days of application. Vigorous crop growth will assist in suppressing weeds not completely killed and those germinating later.

MIXING

TITAN Immense Herbicide is a water-soluble liquid (SL) formulation. Pour the required amount of the product into a spray tank containing almost the total amount of water required. Mix thoroughly. If TITAN Immense Herbicide is added during filling, foaming may occur. If excessive foaming becomes a problem, add a silicone based antifoaming agent at the manufacturers recommended rate. DO NOT use a suction probe unless the antifoaming agent has already been added to the spray tank water. Consult your distributor for specific information on suitable antifoaming agents.

When tank mixing this product with other recommended compatible products, first add the other product(s) to the tank and mix thoroughly before adding TITAN Immense Herbicide.

TITAN Immense Herbicide may be applied in hard or soft water.

The product is corrosive to mild steel. Use ONLY stainless steel, fibreglass, plastic or plastic-lined containers for mixing, storage and application.

APPLICATION

DO NOT apply by aircraft.

Apply in minimum 70L water per hectare. When the crop is very leafy or when the total weed population exceeds 200 plants/m², apply in a minimum of 100L water per hectare to improve contact and coverage. TITAN Immense Herbicide should be applied a minimum of two hours before rainfall or irrigation. If tank-mixed with other products, follow recommendations for the mixing partner should these extend beyond two hours.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of TITAN Immense Herbicide and before use with other products. If tank-mixed with other products, also follow clean-up procedures recommended for the mixing partner.

COMPATIBILITY

TITAN Immense Herbicide is chemically compatible with TITAN Alpha-Cypermethrin 100 Insecticide and Omethoate 290 Insecticide. It is also chemically and biologically compatible with TITAN Clopyralid 750 SG Herbicide, TITAN LVE MCPA 570 Herbicide, TITAN MCPA 750 Selective Herbicide, TITAN Bromoxynil 200 Selective Herbicide, TITAN Bromoxynil MA Herbicide and TITAN Tralkoxydim 400 WG Herbicide and chemically compatible with TITAN Clethodim 240 Herbicide.

DO NOT tank mix with foliar fertilisers.

All tank mixes are chemically stable over a 24-hour period. In the event of delayed spraying, store tank load out of direct sunlight and maintain agitation if possible. Mixes with TITAN Clopyralid 750 SG Herbicide are also biologically stable over a 24-hour delay. Biological stability of other mixes is unknown.

Mixes with Lontrel Advanced Herbicide in imidazolinone-tolerant Wheat and imidazolinone-tolerant Barley: Tank mixes with Lontrel Advanced Herbicide at 150mL/ha will provide control of composite and legume weeds. Refer to Lontrel Advanced Herbicide label.

Mixes with 570g/L MCPA formulations, eg TITAN LVE MCPA 570 Herbicide in imidazolinone-tolerant Wheat and imidazolinone-tolerant Barley: Tank mixes with 570g/L MCPA formulations, eg TITAN LVE MCPA 570 Herbicide at 500mL/ha will provide control of composite and brassicaceous weeds. Refer to TITAN LVE MCPA 570 Herbicide label.

Mixes with Lontrel Advanced Herbicide in imidazolinone-tolerant Canola: Lontrel Advanced Herbicide aids in the control of legume and composite weed

species, such as Annual Medics (*Medicago* spp.), Capeweed (*Arctotheca calendula*), Chickpea (*Cicer arietinum*), Faba Bean (*Vicia faba*), Field Pea (*Pisum sativum*), Lentil (*Lens culinaris*), Narrow Leaf Lupin (*Lupinus angustifolius*). Use rates above 150mL/ha when these weeds are primary weeds in the paddock and when required by their growth stage. Lontrel Advanced Herbicide above 150mL/ha can slightly impair grass control.

For Chickpea, Faba Bean, Lentil: If targeting Chickpeas and Lentils up to 6 leaf stage and Faba Beans up to 4 leaf stage, use a tank mix of TITAN Immense Herbicide with 250mL/ha Lontrel Advanced Herbicide. The addition of Lontrel Advanced Herbicide does not affect the control of other weeds controlled by TITAN Immense Herbicide. Refer to the Lontrel Advanced Herbicide label.

BEST MANAGEMENT PRACTICE (BMP) PROGRAM

A detailed program has been developed that outlines sound agronomic and integrated weed management practices, designed to optimise the performance of imidazolinone-tolerance systems and minimise the potential for the development of herbicide resistance in weed populations. Consultation on BMP should be undertaken with an imidazolinone-tolerant accredited agronomist prior to use of TITAN Immense Herbicide in the imidazolinone-tolerance systems for wheat, barley and canola. Implementation of the BMP is an essential part of herbicide resistance management.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth of imidazolinone-tolerant Wheat, imidazolinone-tolerant Barley and imidazolinone-tolerant Canola but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under poor growth conditions. The imidazolinone-tolerant Wheat varieties CL STL and CL JNZ have limited tolerance to TITAN Immense Herbicide.

Application of TITAN Immense Herbicide to these Wheat varieties can cause unacceptable injury especially under cold, wet conditions. TITAN Immense Herbicide should not be used on these varieties. DO NOT use this product on any Canola variety other than certified varieties with imidazolinone-tolerance technology. Extreme crop damage and/or death will result to conventional and other herbicide tolerant Wheat and Canola varieties.

FOLLOWING CROPS

This product is broken down in the soil by microbes in wet, aerobic conditions. Under conditions that do not favour breakdown, carry-over soil residues can affect susceptible follow crops. Normally safe residue levels may still affect follow crops when soil nutrition is low or marginal, when cold and very wet soil conditions prevail, or when soil pathogens or nematodes are present. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible.

Note: When the intention is to grow cereals on imidazolinone-tolerant Canola stubble (treated with TITAN Immense Herbicide) self-sown Canola volunteers must be removed before they mature beyond 2-leaf, all macro and micro-nutrients must be maintained at levels necessary to grow the planned crops and sulfonylureas must not be used.

The following minimum re-cropping intervals (months after application) should be observed:

Months after Application	Follow Crops
0	Imidazolinone-tolerant Wheat, imidazolinone-tolerant Barley, Imidazolinone-tolerant Canola
10	Chickpeas, Faba Beans, Field Peas, Lucerne, Lupins, Pasture Legumes, Vetch, Oats*, triticale*, non-imidazolinone-tolerant Barley, non-imidazolinone-tolerant Wheat
34	Conventional and other herbicide tolerant Canola, all other crops

* Non-imidazolinone-tolerant Barley, Oats, Triticale, non-imidazolinone-tolerant Wheat.

The following additional requirements apply if it is intended to sow these cereals during the next winter season:

- DO NOT apply TITAN Immense Herbicide later than the end of August (no later than the end of July in WA).
- DO NOT use TITAN Immense Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 150mm (for 300-375mL/ha use), 200mm (for up to 500mL/ha use) and 250mm (for 600-750mL/ha use).
- DO NOT use above 375mL/ha in the Lower Great Southern Region of Western Australia.

For all situations, if expected rainfall is not received following use of TITAN Immense Herbicide, consult your local TITAN AG representative before planting



non-imidazolinone-tolerant cereals. In calculating rainfall actually received, place greater emphasis on rain received from application up to the end of spring and lesser emphasis on break rains. If single isolated heavy summer and autumn falls and break rains are required to achieve rainfall targets, it may not be safe to sow non-imidazolinone-tolerant cereals within 10 months of application. Consult your local TITAN AG representative for advice.

RESISTANT WEEDS WARNING

TITAN Immense Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors of acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group 2 herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group 2 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from failure of this product to control resistant weeds.

RE-ENTRY

DO NOT re-enter treated areas until spray has dried. If re-entry is necessary wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

DO NOT spray within 50m of wetlands or waterways.

PROTECTION TO LIVESTOCK

TITAN Immense Herbicide is of low hazard to bees.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

1L-100L: KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, cool well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

110L & 1000L: KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the containers and preparing product for use, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and a washable hat and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing. When tank mixing with other products consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Immense Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Toxic to aquatic life with long lasting effects. Precautionary Statements: Avoid release to the environment. Collect spillage. Dispose of contents/ container in accordance with local/regional/national regulations.

† Other Trademarks

