

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

MCPA 750

SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 750g/L MCPA present as the DIMETHYLAMINE SALT

GROUP **4** HERBICIDE

For the control of broadleaf weeds in Cereals, Linseed, Poppies, Pastures, Rice, Sugarcane and Turf
as specified in the Directions For Use table.

APVMA Approval No.: 64337/149436

Pack Size: 20L-1000L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | titanag.com.au

AC

Formulation Type
Aqueous Concentrate

IN A TRANSPORT EMERGENCY
• DIAL 000 • POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING NOT
A DANGEROUS GOOD ACCORDING
TO THE AUSTRALIAN DANGEROUS
GOODS (ADG) CODE FOR TRANSPORT
BY ROAD AND RAIL

DIRECTIONS FOR USE

Restraints: DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

DO NOT apply if rain is likely within 6 hours.

SPRAY DRIFT RESTRAINTS

DO NOT apply by a vertical sprayer.

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plan or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

1. FIELD CROPS				
SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Wheat, Barley, Oats, Triticale	Refer to Weed Table	TAS, SA only	460mL-1.45L	Apply through the tillering stage after the crop plants have 5 leaves and before they commence jointing.
		QLD only		Apply from early tillering (4 to 5 fully emerged main stem leaves plus one or more tillers) until before the boot stage (visible swelling of the head at the top of the main stem).
		NSW, ACT only		Apply 460mL rate after crop has reached 5 leaf stage and weeds marked # in weed table have 3-6 leaves. Higher rates after the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller.
		VIC only	460-960mL	Apply 460mL rate after crop has reached 2-3 leaf stage and weeds marked # in weed table have 2-4 leaves. Higher rate after the crop plants have 5 leaves to fully tillered.
		WA only	600mL-1.35L	Apply to crop from 5 leaf to flag leaf emergence (Zadoks 15-37).
Wheat, Barley, Triticale, Cereal Rye	Capeweed, Double Gee, Erodium, Mustard, Rapistrum, Wild Turnip, Wild Radish	WA only	270mL plus 350mL diuron (500g/L flowable)	Crop: 3-4 leaf stage (Zadoks 13-14).
			330mL plus 500mL diuron (500g/L flowable)	Crop: 4-5 leaf stage (Zadoks 14-15). Apply by aircraft or boom with 20-100L/ha water, when most weeds have germinated and are in 2-5 leaf stage. Crop and weeds should be dry at time of application. Some temporary yellowing of crop may occur after application. Undersown sub-clovers may be slightly retarded. DO NOT apply to undersown Medics.
	Amsinckia (Yellow Burr), Capeweed, Charlock Deadnettle, Double Gee (Spiny Emex, Three-cornered Jack), <i>Erodium</i> spp., Mustard, Turnip Weed, White Ironweed (Corn Gromwell, Sheep Weed), Wild Radish, Wild Turnip	NSW, ACT, VIC only	330mL plus 500mL diuron (500g/L flowable)	Apply by boom spray or aircraft when the crop is at the 3-5 leaf stage (NSW, ACT and SA only) and 5 leaf stage (VIC only). Weeds must be in the 2-5 leaf stage. DO NOT apply to undersown Medic or Lucerne. DO NOT use on Kamilaroi, Kite, Olympic or Shortim Wheat.
	Capeweed, Mustard, Three-cornered Jack, Turnip Weed, Wild Radish, Wild Turnip, Wards Weed, Common Ice Plant, Pimpernel	SA only	230mL plus 500mL diuron (500g/L flowable)	DO NOT use on very light sandy soils.
Wheat, Barley, Oats, Triticale	Soldier Thistle	SA only	460mL plus 200mL TITAN Clopyralid 300	Apply to crop at tillering stage after the crop has 5 leaves and before the crop commences jointing.
	Skeleton Weed		660mL plus 500mL TITAN Clopyralid 300	Apply to rosettes from 5-15cm diameter when the crop is between the leaf and late tillering growth stages. It is essential that all the Skeleton Weed is emerged and minimum of 5cm diameter which is usually not before August. This rate will provide control of Skeleton Weed until harvest and will substantially reduce seed numbers the following season.

1. FIELD CROPS – continued				
SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Field Peas	Prickly Lettuce	NSW, VIC, TAS, SA only	Weed Stage: 4 leaf, 80mL plus 125mL diflufenican (500g/L)	Field Peas: Apply early post-emergence after the third node stage and before the start of flowering. Warning: Field Peas grown on high pH soils in the presence of free lime may be less tolerant to diflufenican (500g/L). Application and Weed Control: Apply when weeds are actively growing. For optimum results apply 4 to 6 weeks post-sowing. Application beyond 8 weeks post-sowing may result in reduced levels of weed control. In most situations the rate specified for each weed size will give satisfactory control. Under certain conditions such as: • High crop and weed density, • Late season germinations, • Abnormal weed growth (including early flowering); higher rates of product (up to the maximum rate of application specified for that weed) may be required. Diflufenican (500g/L) will not effectively control: • Regrowth of suppressed weeds, • Transplanted weeds, • Regrowth from rhizomes or roots, • Weeds growing under stress from previous herbicide applications.
			Weed Stage: 6 leaf, 100mL plus 150mL diflufenican (500g/L)	
	Charlock (Wild Mustard)	NSW, VIC, TAS, SA only	Weed Stage: 4 leaf, 80mL plus 125mL diflufenican (500g/L)	
			Weed Stage: 6 leaf, 100mL plus 150mL diflufenican (500g/L)	
Linseed	Refer to Weed Table	QLD only	460mL-1.45L	Apply when crop is 10-15cm tall with at least 170L/ha of water. DO NOT spray after buds appear. Some wilting effect may be seen following application but crop usually recovers quickly. DO NOT apply by aircraft.
		NSW, ACT, VIC, SA, TAS only	460-730mL	
Oilseed Poppies (<i>Papaver somniferum</i>)	Fat Hen, Spear Thistle	NSW, TAS only	1.0L	Use only in consultation with the poppy contracting company. Apply at least one week after petal fall after poppy seed has been set. Apply by aircraft in 30-40L of water/ha. This treatment will control late germinating weeds if they are green and at the bud or early flowering state. Spraying at this time will twist weeds below the crop canopy. DO NOT apply in horticultural areas where there is a risk of spray or vapour drift over susceptible crops.
Sugarcane	Blue Top, Chinese Burr, Flannel Weed, Gambia Pea, Bell Vine, Streaked Rattle Pod, Bindweed, Pink Convolvulus, Cupids Flower, Merremia Vine, Morning Glory	QLD only	930mL	Post-emergent: Apply as a directed inter-row spray. DO NOT exceed rates with phenoxy sensitive varieties.
	As above plus Fat Hen, Noogoora Burr		1.45L	
Rice	Dirty Dora, Docks, Sedges, Starfruit, Buddah (Butter) Pea	ALL STATES	485mL to 970mL	Apply by aircraft at early/mid-tillering stage of Rice but not later than panicle initiation. Use 485mL/ha at 35 days after aerial sowing (approx. 2 tillers per Rice plant). Use 970mL/ha at 45 days or more after cereal sowing (3 tillers or more per Rice plant). The crop should be sprayed before flowering of the weeds. Water Management: Before herbicide application, lower water level in the bays to at least two thirds of the weed growth for direct contact with MCPA 750. Weeds covered by water will not be controlled. Raise water depth to normal levels 24 hours following spraying.
Flax	Refer to Weed Table	VIC, TAS only	325mL	Apply to crop when 8-20cm high.
2. PASTURES, SEED CROPS & TURF				
SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Grass Pastures	Refer to Weed Table	ALL STATES	460mL-2.7L	Apply to established grass based pastures. Damage may occur to legumes if present. See "Legume Tolerance" under General Instructions. Note: Capeweed, Paterson's Curse and Variegated Thistle may be poisonous to stock after spraying.
	Salvation Jane	SA only	660mL plus 400mL diuron (500g/L flowable)	Use this mixture only if spraying is delayed until early winter. Rosettes should not exceed 20cm diameter.

2. PASTURES, SEED CROPS & TURF – continued				
SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Grass Pastures – continued	Soldier Thistle	SA only	460mL-1.45L	Use low rate on seedlings with no more than 2 true leaves. Use 100 litres of water per ha for best results. If clover present do not exceed 575mL.
	Seedling and established Horehound	NSW, ACT only	2.0L	Apply in autumn when Horehound plants are at the seedling stage. If Sub-Clover or Medic plants are present they will be damaged by this application.
Grass Seed Crops	Refer to Weed Table	QLD, VIC, TAS, SA, WA only	460mL-2.7L	Apply to established grass seed crops.
Sub-Clover Seed Crops	Refer to Weed Table	VIC, SA only	460-960mL	DO NOT exceed rate specified. Spray in autumn and early winter.
Pastures – Spray/Graze Technique	Caltrop, Capeweed, Charlock, Mustards, Paterson’s Curse, Shepherd’s Purse, Saffron, Slender Spear or Variegated Thistle, Turnip Weed, Wild Radish, Wild Turnip	NSW, ACT, VIC only	230-930mL	Spray actively growing 6-8 weeks old weeds. Introduce stock 7-10 days after spraying, preferably sheep as cattle are less effective. Use stocking rate at least 5 times heavier than normal until weeds have been reduced but before survival of desirable pasture species is threatened. Lucerne and Medics may be damaged and should be grazed short before spraying. See “Legume Tolerance” under General Instructions.
		TAS only	460mL	
	Refer to Weed Table	WA only	610mL-1.25L	
	Capeweed, Dandelion, seedling Dock, Three-cornered Jack, Geranium, Erodium, Mustard, Annual Thistles, Turnip Weed, Wild Radish, Wild Turnip	SA only	330mL	Precaution: An increased quantity of poisonous plants may be eaten by stock using spray/graze eg. Caltrop, Capeweed, Paterson’s Curse and Variegated Thistle and deaths could result from causes such as nitrate poisoning. With Paterson’s Curse, preferably graze stock soon destined for slaughter and avoid extended periods of grazing. Avoid grazing with young or breeding stock. DO NOT graze horses or pigs on Paterson’s Curse. Observe withholding period of at least 7 days after spraying.
	Soldier Thistle		395mL	
Turf	Refer to Weed Table	ALL STATES	330mL or 2.8mL per 100 sq. metres	Newly laid: Delay until well established or at least 2 months old. Apply in high volume to actively growing weeds. See “Legume Tolerance” under General Instructions. Buffalo Grass is more easily damaged than other grasses.
			930mL-1.8L or 10-19mL per 100 sq. metres	Established: Apply in high volume to actively growing weeds. DO NOT mow for 2 days before application. Some transitory damage may occur to fine turf grasses and Strawberry Clover. See “Legume Tolerance” under General Instructions. Buffalo Grass is more easily damaged than other grasses.
3. SPOT SPRAYING				
SITUATION & CROP	WEEDS CONTROLLED	STATE	MIXING RATES/COMMENTS	
High Volume Spraying	Refer to Weed Table	ALL STATES	Use 130mL/100L of water for easy-to-kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 270mL/100L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.	
Knapsack Application	Refer to Weed Table	ALL STATES	Use 13mL/10L of water for easy-to-kill weeds such as Wild Turnip, Mustards and Wild Radish. Use 27mL/10L of water for more difficult to control weeds such as Docks, Hoary Cress and Bindweed.	
4. NON-CROP AREAS, FALLOW LAND, INDUSTRIAL & COMMERCIAL AREAS, RIGHTS-OF-WAY				
SITUATION & CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Non-Crop areas, Fallow Land, Industrial and Commercial Areas, Rights-of-way	Refer to Weed Table	ALL STATES	Refer to Weed Table	Apply at rates shown in Weed Table when weeds only are present and non-crop or pasture is involved.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

Cereal Grain: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

Pasture: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

Poppies: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

Harvest Withholding Period (other than Poppies and Cereal Grain): NOT REQUIRED WHEN USED AS DIRECTED.

WEED TABLE: Where weeds are to be sprayed in a crop or pasture situation, use only the rate given for the particular crop or situation indicated under the Directions For Use table. In most cases this will give control, however some hard-to-kill weeds or those in advanced stages of growth may only be suppressed. The rates listed in the Weed Table below are for use where weeds only are present and no crop or pasture is involved. Use at these rates in a crop or pasture other than for spot spraying may cause damage. This product should only be used in those States where a rate or range of rates is indicated for the particular weeds listed.

WEEDS CONTROLLED	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS
	VIC	NSW, ACT	SA	TAS	QLD	WA	
Ball Mustard	-	-	1.15L	-	-	-	Apply at rosette stage.
Bathurst Burr	1.35L	660mL-1.35L	1.35L	1.35L	1.35L	1.35L	Spray young seedlings only.
Black Bindweed	-	970mL-1.35L	-	-	930mL-1.35L	-	Apply at young stage prior to flowering.
California Burr	-	660mL-1.35L	-	1.45L	-	-	Spray young seedlings only.
Cape Tulip	-	-	-	-	-	1.35L	Apply just prior to flowering. Gives mild suppression only.
Capeweed	-	1.45L	1.45L	1.45L	1.45L	-	Apply at seedling stage.
Carrot Weed	1.35L	-	-	-	1.35L	-	Apply at seedling stage.
Charlock	460mL	660mL	460mL	660mL-1.45L	460mL	-	Spray up to rosette stage.
Cutleaf Mignonette	-	-	1.45L	-	-	-	Apply at young rosette stage.
Dandelion	1.35L	1.35L	1.35L	1.35L	1.35L	1.35L	Apply at young rosette stage.
Deadnettle	-	1.45L	-	-	1.45L	-	Moderately susceptible. Apply at seedling stage.
Docks	960mL	-	1.45L	1.45L	-	-	Apply at rosette stage and before flower heads emerge. Only the top growth of old established plants will be affected.
Fat Hen	1.35L	660mL-1.35L	-	1.35L	1.35L	-	Spray up to rosette stage.
Fennel	1.80L	1.80L	1.80L	1.80L	1.80L	-	Apply at young stage.
Field Bindweed	1.35L	-	1.35L	1.35L	1.35L	1.35L	Apply at bud stage, repeating over several seasons.
Field Cress	-	-	-	1.45L	-	-	Apply at young stage.
Flatweed or Catsear	-	-	-	1.45L	-	930mL	Apply when weeds are 10-15cm high.
Fumitory (red)	930mL	930mL	930mL	-	930mL	-	Apply at young stage prior to flowering.
Hoary Cress	3.0L	-	1.45L	-	3.0L	-	Apply before flowering.
Horehound	-	2.0L	1.45L	-	-	-	Apply at rosette stage.
Hedge Mustard	730mL [#]	-	460mL	460mL	460mL	825mL	Apply at rosette stage.
Hexham Scent or Melilotus	960mL	800mL	930mL	-	930mL	-	Apply at young seedling stage.
Lesser Swinecress	-	-	-	1.45L	-	-	Apply at young stage.
Lincoln Weed	-	-	460mL	-	-	-	Apply at rosette stage.
London Rocket	-	-	-	-	-	825mL	
Lupins	-	460-960mL	-	-	-	825mL	Spray up to 10cm high.
Mintweed	1.35L	1.35L	-	-	1.35L	-	Spray young seedlings only.
Mustards	730mL [#]	660mL [#]	730mL	730mL	-	825mL	Spray up to rosette stage.
Noogoora Burr	1.35L	660mL-1.35L	1.35L	-	1.35L	1.35L	Spray young seedlings only.
Opium Poppy	-	-	-	1.45L	-	-	Apply at young stage.
Paterson's Curse	-	660-960mL	-	-	2.70L	-	Apply early rosette stage.
Pimpernel	-	-	460mL	-	-	-	Apply at rosette stage.
Plantains	-	1.35L	-	1.35L	1.35L	-	Apply at young stage.
Rapeseed	-	660-960mL	-	-	-	-	Spray up to rosette stage.
Rough Poppy	-	460-960mL	-	-	-	-	Spray up to rosette stage.
Safflower	-	460-660mL	-	-	-	-	Spray up to rosette stage
Shepherd's Purse	-	-	-	1.45L	-	-	Apply at rosette stage.
Skeleton Weed	-	960mL-1.35L	-	-	-	-	Spray rosettes before aerial growth commences.
Sorrel	-	-	1.45L	-	-	-	Apply to rosette stage.
Stinkwort	1.35L	660-960mL	1.35L	-	-	1.35L	Apply when weeds are 10-15cm high.
Sunflower (seedlings)	-	690mL-1L	-	-	-	-	Spray up to rosette stage.
Thistles:							
- Artichoke	1.05L	-	1.45L	-	1.35L	460mL	Apply at rosette stage.
- Nodding	-	1.65L	-	-	-	-	Spray early rosettes. Repeat spray required in following spring or autumn.
- Saffron	960mL-1.65L	660mL-1.35L	930mL-1.65L	930mL-1.65L	930mL-1.65L	960mL-1.65L	Spray up to rosette stage (September – WA).

WEEDS CONTROLLED	APPLICATION RATE PER HECTARE						CRITICAL COMMENTS
	VIC	NSW, ACT	SA	TAS	QLD	WA	
Thistles continued:							
- Scotch or Cotton	1.45L	-	460-930mL	1.45L	-	-	Apply at seedling stage.
- Slender	2.0L	660mL-1.35L	660mL	2.0L	-	2.0L	Spray young seedlings only.
- Soldier	730mL	-	730mL	-	-	-	Spray young seedlings. Use higher rate for larger weeds.
- Spear	-	960mL-1.35L	-	1.45L	-	-	Apply at rosette stage.
- Star	-	-	730-930mL	-	-	-	Apply at seeding stage. Use higher rate for larger weeds.
- Variegated	960mL	660mL-1.35L	460-930mL	960mL	960mL	960mL	Spray young seedlings only. Use 1.35L/ha (SA) for well-developed weeds.
Turnip Weed	730mL [#]	660mL [#]	660mL	-	-	-	Spray up to rosette stage.
Wild Artichoke	-	-	1.45L	-	-	-	Apply at rosette stage.
Wild Radish	930mL	660mL [#]	930mL	1.35L	460-730mL	800mL	Spray up to rosette stage.
Wild Turnip	730mL	660mL [#]	460mL	1.35L	460-730mL	600mL	Spray up to rosette stage.

GENERAL INSTRUCTIONS

Weeds should be sprayed while actively growing and at their most susceptible stage. As a general guide, annuals should be sprayed when young and perennials just prior to appearance of bud stage. Extremes of cold or drought are unfavourable. The best conditions are when the soil is moist, weather fine and rain unlikely within 6 hours following application.

DO NOT spray crops or weeds outside the stages indicated in 'Critical Comments' as damage, loss of yield or inadequate weed control may result. DO NOT spray in high winds.

APPLICATION/MIXING INFORMATION

This product mixes readily with water. Half fill the spray tank with clean water and add the required amount of product. Agitate thoroughly before adding the remaining water.

BOOM SPRAYING

- Use 30-120 litres of water per hectare.
- Use minimum of 170 litres water per hectare for Linseed.

AERIAL SPRAYING

- Use 10-90 litres water per hectare.

EQUIPMENT MAINTENANCE AND USAGE

Equipment that has been used for this chemical should not be used for the application of other materials to sensitive plants unless it has been cleaned with Spraymate Tank and Equipment Cleaner or well washed out with a 2% solution of cloudy ammonia followed by several clear water rinses.

COMPATIBILITY/TANK MIXES

This product can be tank mixed with TITAN Metsulfuron 600, Banvel*, TITAN Bromoxynil 200, Flowable Diuron, TITAN Clopyralid 300, Sertin*, TITAN Paraquat 250, Flowable Terbutryn, Tordon* and Tribunil*. The effectiveness of Imidan*, Le Mat*, Metasystox* and dimethoate may be reduced by alkaline hydrolysis if tank mixed with this product. For these situations the addition of Spraybuff* or Spraymate* L1-700 is recommended. When tank mixing add the required amount of compatible product (and surfactant if recommended) when the spray tank is nearly full and while continuously agitating. If left overnight, agitate thoroughly before spraying.

RESISTANT WEEDS WARNING

TITAN MCPA 750 Selective Herbicide is a member of the Phenoxy group of herbicides. This product has the disruptors of plant cell growth mode of action. For weed resistance management this product is a Group 4 Herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Drift Warning: DO NOT use on or in situations where damage to susceptible crops – such as cotton, vines, tomatoes, vegetables, fruit trees, lucerne, legume crops and pastures, oilseed crops or other susceptible crops and trees (eg. Kurrajongs, Belahs, Eucalypts) – may occur from direct application, spray drift or vapours.

LEGUME TOLERANCE

The use of MCPA is preferred to 2,4-D Amine (Amicide) or Ester (Estericide) in situations where Legumes are undersown in the crop.

Note: The tolerance of Pasture Legumes to applied product can vary with species, variety, growing conditions, stage of growth and companion crop cover. DO NOT spray cereal crops undersown with Lucerne as damage may result.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and goggles. If product is on skin, immediately wash area with soap and water. If product or spray is in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN MCPA 750 Selective Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if swallowed. Causes skin irritation. Causes serious eye damage. Very toxic to aquatic life with long lasting effects. Precautionary Statements: Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves / eye protection / face protection. IF SWALLOWED: Call a POISON CENTER/ doctor if you feel unwell. IF ON SKIN: Wash with plenty of water. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER/doctor. Specific treatment (see on this label). Rinse mouth. Take off contaminated clothing and wash it before reuse. If skin irritation occurs: Get medical advice/attention. Collect spillage. Dispose of contents/container in accordance with local/regional/national/international regulations.

