

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

TITAN

PICTRON EC

HERBICIDE

**ACTIVE CONSTITUENTS: 300g/L TRICLOPYR present as the BUTOXYETHYL ESTER
100g/L PICLORAM present as HEXYLOXYPROPYLAMINE SALT**

GROUP 4 HERBICIDE

For control of a range of environmental and noxious woody and herbaceous weeds as specified in the
Directions For Use table.

APVMA Approval No.: 62484/140065

Pack Size: 1L-20L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | Fax (02) 9999 0483
titanag.com.au

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL

DIRECTIONS FOR USE

Restraints:

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, damage or previous herbicide treatment, as reduced levels of control may result.

DO NOT spray if rain is likely within one hour or if foliage is wet from rain or dew.

DO NOT burn off, cut or clear blackberry or other woody weeds for at least six months after spraying.

DO NOT apply by aerial application in wind in excess of 15km/hr and air temperatures above 35°C.

In areas prone to flooding, treatment should commence after any annual flooding as such areas flooded within 9 months following application may have reduced results.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at www.apvma.gov.au/spraydrift

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

1. WOODY WEED SITUATIONS

Table A: High Volume Spraying – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/100L	CRITICAL COMMENTS
African Boxthorn	Less than 2m tall	ALL STATES	500mL	Apply when bushes have good leaf cover, growth and no leaf fall.
<i>Angophora</i> spp.	1 to 3m tall	NSW, ACT only	350mL	
Australian Blackthorn	Less than 2m tall		500mL	Apply from late spring to early autumn.
<i>Banksia</i> spp.	1 to 3m tall		350mL	
Biddy Bush (Chinese Shrub) (Sifton Bush)	Autumn when actively growing		500mL	Add TITAN Wetter 1000 Wetting Agent at 125mL/100L of water for best results.
Blackberry in association with: Docks, Ragwort, Smartweed Thistles	Late spring to autumn	ALL STATES except NT	350mL or 500mL	Use the higher rate on plants that have been damaged by grazing stock or insects and on known difficult to kill Blackberry. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with TITAN Pictron EC Herbicide.
Blue Heliotrope	Flowering	QLD, NSW only	500mL	Apply in a minimum spray volume of 1250L/ha.
Brooms: Cape, English, Flax Leaf, Montpellier	Spring to mid-summer prior to pod formation	ALL STATES except NT	250mL	Apply as thorough foliage spray.
	Autumn to winter		350mL	
Camphor Laurel	Less than 2m tall	ALL STATES	500mL	
	Above 2m tall			
<i>Casuarina</i> spp.	1 to 3m tall		350mL	
Chinese Apple	Less than 2m tall	QLD, WA only	350mL	Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Cockspur Thorn	Spring to autumn	QLD, NSW, ACT only		Apply as a thorough foliage spray.
Common Sensitive Plant	Any time when actively growing	QLD, WA, NT only	200mL	To avoid leaves closing during application, spray plants while moving forward. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Crofton Weed	Spring to autumn	QLD, NSW, ACT only	350mL	Apply as a thorough foliage spray.
Eastern Cotton Bush (<i>Maireana microphylla</i>)	Spring to autumn	QLD, NSW only	500mL	Add Uptake Spraying Oil at 500mL/100L of water. Some bushes may require a follow-up spray to control regrowth.
<i>Eucalyptus</i> spp.	Seedling and regrowth from small lignotubers, 1 to 3m tall	ALL STATES	350mL or 500mL	Apply the high rate where difficult to control species of Eucalyptus regrowth is present. Addition of an adjuvant may improve results, contact TITAN AG for details.
Galenia	Fresh growth in spring to summer	NSW only	500mL	Use 2000L of spray mixture/ha.
Giant Bramble	Spring to autumn	QLD, WA, NT only		Penetration of thick clumps may be difficult and respraying may be necessary. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Gorse	1 to 1.5m tall	ALL STATES except NT	250mL	Spring and summer treatment only. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
	Over 1.5m tall or autumn treatment		350mL	Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
	Winter treatment		500mL	Brownout may not be complete until summer. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.

1. WOODY WEED SITUATIONS – continued				
Table A: High Volume Spraying – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.				
AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/100L	CRITICAL COMMENTS
Groundsel Bush (<i>Baecharis halimifolia</i>)	1 to 1.5m tall in spring and summer	ALL STATES except NT	250mL	Apply as a thorough foliar spray.
	Over 1.5m tall or autumn treatment		350mL	
Green Cestrum	Late spring to early autumn	QLD, NSW, ACT only	500mL	One application may give satisfactory control. Any subsequent regrowth and seedlings must be resprayed at approximately 1m high.
Hawthorn	Less than 2m tall	ALL STATES	500mL	Apply from late spring to early autumn.
Horehound	Pre-flowering		350mL	Apply as a thorough foliar spray.
Japanese Sunflower	Pre-flowering	QLD, NSW only		
Lantana (<i>Lantana camera</i> , <i>Lantana montevidensis</i>)	Up to 1m tall in summer to autumn	ALL STATES		Add one of the following adjuvants when using 350mL rate: Uptake Spraying Oil @ 0.5% v/v, TITAN Organosilicone Surfactant.
	1 to 2m tall in summer to autumn		500 or 750mL	
Lion Tail (<i>Leonatis nepetifolia</i>)	Pre-flowering	QLD only	200mL	Apply as a thorough foliar spray. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Limebush	Any time of year with good leaf cover and soil moisture	QLD, NT only	350mL	Penetration of thick clumps may be difficult and respraying may be required. Addition of an adjuvant may improve results – contact TITAN AG for details.
Manuka	At flowering	VIC only	500mL	For optimum results, add TITAN Organosilicone Surfactant at 200mL/100L of spray. Thoroughly wet foliage, stems and soil around the base of the plants.
Mesquite (<i>Prosopis</i> spp.) (<i>Prosopis velutina</i>)	Seedling, full leaf and flowering before podding	QLD, NSW, WA, NT only	350mL	DO NOT spray plants bearing pods. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
		QLD only	670mL	
Mistflower	Spring to autumn	QLD, NSW, ACT only		Apply as a thorough foliar spray.
Mother-of-millions	Flowering	QLD, NSW only	500mL	Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Paddy's Lucerne	Active growth	NSW only	500mL	Plants that have been continually slashed or grazed over many seasons may be difficult to control and regrowth may occur.
Parkinsonia	Under 2m tall	QLD, WA, NT only	350mL	Add Uptake Spraying Oil at 500mL/100mL water. Avoid spraying under dry conditions when plants are stressed or bearing pods. Thoroughly wet foliage.
Prickly Pear (common), Smooth Tree Pear	Active phyllode growth	ALL STATES	500mL	Apply as a thorough foliage spray. Regrowth may occur, so a follow-up application may be necessary.
Rubber Vine (Not infected with rust)	Up to 1.5m tall at flowering	QLD, WA, NT only	350mL	Spray all leaves and stems just to the point at run-off and thoroughly spray the base of the plant. With larger more dense stands regrowth may occur. Subsequent control of any regrowth should be done by basal bark spraying.
	Dense stands greater than 1.5m tall at flowering		500mL	
Siam Weed	Active growth	QLD, WA only	350mL	Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Sicklepod	Up to flowering	QLD, WA, NT only	200mL	DO NOT apply to podding plants. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
St John's Wort	From flowering to early seed set	ALL STATES	500mL	Late spring to early summer.
Sweet Briar	Up to 1.5m tall	ALL STATES except NT	350mL	Add metsulfuron-methyl at 10g/100L water to obtain more reliable results with the lower rate of TITAN Pictron EC Herbicide.
			500mL	Full leaf to ripe fruit prior to leaf fall. Thorough wetting including the crown is recommended.
Tobacco Weed	Actively growing plants	QLD, WA, NT only	300mL	Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
Wattle (<i>Acacia</i> spp.) (except Corkwood Wattle)	1 to 3m tall	ALL STATES	350mL	
Wild Rosemary (<i>Cassinia laevis</i>)	Active growth, 0.5 to 1m tall	QLD only	350 to 500mL	Use lower rate on seedlings 0.5m tall. Apply as a thorough foliar spray.
Wild Tobacco Tree	Spring to autumn up to 2m tall	QLD, NSW, ACT only	350mL	

1. WOODY WEED SITUATIONS – continued				
Table B: Aerial Application – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.				
AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry	Summer to autumn	QLD, NSW, VIC, SA, WA only	10L	Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with TITAN Pictron EC Herbicide. Warning: Eucalyptus species up to 4m may be killed if sprayed during this treatment. Mature trees which are 15 to 20m tall may be partially defoliated but are likely to recover.
Gorse		TAS only		Helicopter application only.
Cockspur Thorn, Crofton Weed, Lantana, Mistflower	Late autumn	QLD, NSW, NT only	1.5L plus 7.5L 2,4-D Amine (500g/L)	Spray with calibrated equipment using the half overlap opposite pass technique applying a minimum spray volume of 150L/ha. Follow up respraying will be required.
Lantana			10L	Helicopter application only.
Rubber Vine (Not infected with rust)	When flowering	QLD, NT only (helicopter only)	3L to 5L	Use rates will depend upon the density and height of the Rubber Vine stand. The higher rates should be used on dense stands, however, complete coverage and penetration may be difficult. Follow up respraying will be required. Any regrowth should be sprayed with a suitable Basal Bark herbicide.
St John's Wort	Flowering to early seed set (Nov-Jan)	NSW only	4L	Helicopter application only. Follow up spraying will be required in the following season.
AGRICULTURAL NON-CROP AREAS ON FLOOD PLAINS				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Parkinsonia	Seedlings, 1-2m tall or 12-24 months old	QLD, NT only (helicopter only)	3L	Add Uptake Spraying Oil at 1L/ha.
Table C: Controlled Droplet Application (CDA) – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.				
AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry In association with: Docks, Ragwort, St John's Wort, Thistles	Summer to autumn	ALL STATES except NT	Apply undiluted	One application may give satisfactory control but subsequent regrowth and seedlings should be resprayed after hardening off. Where herbicides other than Group 4 herbicides have been used, allow two seasons regrowth to occur before respraying with TITAN Pictron EC Herbicide.
Table D: Low Volume High Concentrate Application Techniques – (Gas Powered Gun, Sprinkler Sprayer) – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.				
AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry	Late spring to autumn	QLD, NSW, ACT, TAS, SA, WA only	335mL	Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Camphor Laurel, Cockspur Thorn, Crofton Weed	Less than 1.5m high	QLD, NSW, ACT only	500mL	
Eucalyptus species	Seedlings up to 2m tall	ALL STATES	335mL	
Mistflower	Less than 1.5m high	QLD, NSW, ACT only	500mL	
Sweet Briar	1.5m tall, full leaf to ripe fruit	NSW only		Gas Powered Gun only: Apply to actively growing bushes not more than 1.5m tall that have not more than 5 stems from the crown.
St John's Wort	During flowering to early seed set	NSW, VIC, TAS only		Gas Powered Gun only: One application should provide control. Minor regrowth and seedlings may be retreated the following summer.
Wild Tobacco Tree	Less than 1.5m high	QLD, NSW, ACT only		Apply to actively growing bushes which are able to be sprayed on all sides. For larger bushes, the high volume application technique is recommended.
Table E: Boom Application – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Galenia	Fresh growth during spring to summer	NSW only	5L	Rough mine sites will require adequate spray equipment such as boomless nozzles for effective coverage.
Sicklepod	Up to flowering	QLD, NT only	3L	DO NOT apply to podding plants. Add TITAN Wetter 1000 Wetting Agent at 100mL/100L of water for best results.
St John's Wort	Flowering to early seed set (Nov-Jan)	NSW only	2 to 4L	Use the higher rate on dense infestations and when longer residual control is required. Follow up respraying will be required in the following season.

2. FALLOW SITUATIONS

Table A: Boom Application – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE/ha	CRITICAL COMMENTS
Blackberry Nightshade Suppression only	10 to 25cm tall, prior to flowering	QLD, NSW only	200 to 400mL + 12L TITAN Glyphosate 450 Herbicide + Adjuvant	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing. Use the lower rate on the smaller weeds as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i>). Refer to TITAN Glyphosate 450 label for use of adjuvant. DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions – Minimum Recropping Periods: Black Cracking Clay Soils, NSW & QLD. Dry conditions after application will increase the recropping interval.
Camel Melon, Prickly Paddy Melon, Cucumber Melon (<i>Cucumis melo</i>)	From 2 leaf to 50cm diameter			
Common Sowthistle	From 8 leaf to flowering			
Cow Vine	From 2 to 5 leaf up to 15cm diameter, prior to flowering	QLD, NSW only	200 to 400mL + 12L TITAN Glyphosate 450 Herbicide + Adjuvant	FOR USE BY GROUND EQUIPMENT ONLY. Plants must be actively growing. Use the lower rate on the smaller weeds as specified in the weed growth stage (or up to 5cm diameter for <i>Polymeria pusilla</i>). Refer to TITAN Glyphosate 450 label for use of adjuvant. DO NOT plant susceptible crops for up to nine months after application, as specified in General Instructions – Minimum Recropping Periods: Black Cracking Clay Soils, NSW & QLD. Dry conditions after application will increase the recropping interval.
Lucerne (established)	Active growth, 15 to 25cm high, during spring			
<i>Polymeria pusilla</i>	2 to 12 leaf up to 200m diameter, prior to flowering			

Table B: Blanket Wiper Application – See GENERAL INSTRUCTIONS – APPLICATION section for application method details.

FALLOW				
WEEDS CONTROLLED	WEED GROWTH	STATE	RATE	CRITICAL COMMENTS
Bitterbark (<i>Alstonia constricta</i>)	From summer to end of autumn	QLD only	Blanket wipers: 1:4 (1 part TITAN Pictron EC Herbicide to 4 parts water)	Blanket wipers: For best results apply in autumn to tall (> 60cm) plants using two opposite directional passes (up and back). Follow up “missed” plants with a spot spray application. These will be obvious after 6 weeks. Blanket wiper applications can be made in summer when plants are smaller but follow up spraying may be necessary. DO NOT disturb (cultivate) the treated patches for at least 3 months. Best long term control is achieved when patches are left undisturbed for as long as possible after treatment (at least 6 months).
			Spot spraying “missed” plants: 2% solution for spot spraying (eg. 100mL TITAN Pictron EC Herbicide in 5L water)	Spot spraying “missed” plants: Thoroughly wet all stems and leaves without producing any solution run-off. Avoid any spray reaching the soil surface.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

IN TASMANIA FOR BLACKBERRY: DO NOT TREAT BUSHES CARRYING MATURE OR NEAR MATURE FRUIT.

FOR NATIVE VEGETATION: USE OF TITAN PICTRON EC HERBICIDE ON NATIVE VEGETATION MUST BE DONE IN ACCORDANCE WITH STATE AND/OR LOCAL LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MINIMUM RECROPPING PERIODS – Black Cracking Clay Soils, Northern NSW & QLD				
Table A: Boom Application				
Plant-back periods for crops following the application of TITAN Pictron EC Herbicide for rates up to 600mL/ha.				
Rate mL/ha	200	300	400	600
Crop	Months			
Wheat	2	2	4	4
Barley	2	2	4	4
Canola	2	4	4	4
Faba Bean	4	4	6	6
Chickpea	4	6	6	6
Lucerne	6	9	9	9

These plant-back periods are based on a normal rainfall pattern. During drought conditions (or when rainfall is less than 100mm for a period of 4 months or greater) the plant-back period may be significantly longer.

Table B: Blanket Wiper Application	
Plant-back periods for crops following blanket wiper application.	
Crop	Months
Broadleaf Crops	18
Lucerne	6
	This will allow any potential soil residues to dissipate, if any, and allow effective control.

Note: Before using TITAN Pictron EC Herbicide in tank mixes with other herbicides, check the plant-back information on all product labels. The most residual product, ie. the product with the longest plant-back period, will determine the time between spraying and planting.

COMPATIBILITY

Follow any regional restrictions, and all directions and restrictions on the label, of any chemical mixed with TITAN Pictron EC Herbicide (eg. 2,4-D amine). TITAN Pictron EC Herbicide is compatible with the following herbicides: TITAN 2,4-D Amine products, TITAN Metsulfuron 600 WG Herbicide, TITAN Glyphosate 450 Herbicide, TITAN Fluroxypyr products.

TITAN Pictron EC Herbicide is compatible with the following adjuvants, as per Directions for Use: Uptake Spraying Oil, TITAN Organosilicone Surfactant, non-ionic surfactant (1000g/L) or equivalent.

MIXING

SHAKE WELL BEFORE USE

Mix only with water. Half fill the spray unit with water, and add the required amount of TITAN Pictron EC Herbicide. Add the remaining water with the agitator running. If required, then add spray oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

APPLICATION

1. WOOD WEED SITUATIONS

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to approximately 1m in situations which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

A. High Volume Spraying

Thorough coverage of foliage to the point of run-off is essential, however, avoid excess spraying which is wasteful of chemical.

Hand Gun: Apply the recommended mix to give full coverage of leaves and stems through a No.6 to 8 tip at 700 to 1500kPa (400 to 500kPa for St John's Wort). A spray volume of 3000 to 4000L per infested hectare of 1 to 2 metre high blackberry (30 to 40L/m²) should be used. Use 2000L of spray mixture/ha of Galenia infestation (ie. 20L/100m² infested area).

Knapsack: Apply the recommended spray mix to give full coverage of leaves and stems. The final volume of application should be similar to hand gun. A spray volume of 3 to 4L/10m² infested area should be used. A spray volume of 2L/10m² should be used for an area infested with Galenia.

B. Aerial Application

Apply in 200L of water/ha using an aircraft to apply 100L per pass on a double overlap pattern using nozzle configurations to produce droplets of 250 to 350 micron diameter.

The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.

DO NOT spray when wind exceeds 15km/hr and/or air temperature reaches 35°C.

C. Controlled Droplet Application (CDA)

Results similar to high volume spraying can be obtained using Micron Herbi* or similar equipment. Select a nozzle to give a flow rate of 2m/sec and sweeping action of approximately 1m/sec to ensure a droplet density of 20/cm². Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided by CDA unit.

D. Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of entire bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

Gas Powered Gun: Apply 50mL shots to obtain uniform coverage of 4-5m² of surface area of bush. This relates to 20 droplets/cm² of leaf surface.

Sprinkler Sprayer: This technique involves using a micro sprinkler that is connected to a hollow fibreglass rod attached to a pneumatic knapsack sprayer. Use at low pressures (50 to 200kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

E. Boom Application

Application in a minimum spray volume of 200L/ha for Galenia and St John's Wort and 600L of water/ha for Sicklepod. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

2. FALLOW SITUATIONS

A. Boom Application

Application of TITAN Pictron EC Herbicide in a minimum spray volume of 50L/ha is recommended. Flat fan nozzles are recommended, using pressure in the range of 200-300kPa. Boom height must be set to ensure double overlap of nozzle patterns.

B. Blanket Wiper Application: Blanket needs to be made from durable and wettable material with a rigid backing. Blanket should be rigidly mounted behind motorised vehicle (eg. tractor, 4-wheel drive vehicle) and set low but never touching the ground. The chemical solution should be fed to the blanket at a flow rate sufficient to keep the blanket wet but not dripping. In thick patches the blanket may require more frequent solution recharge (rewetting). Ideally, a scraper bar should be mounted in front of the blanket in order to scrape or damage the bark (but not sever the stems) prior to the blanket wiping the stems. This scraper may be mounted at the front of the vehicle. Two passes (in opposite direction) with the blanket increases the contact with the plant. Ground speeds of 10-15kph are ideal for blanket wiping application.

CLEANING SPRAY EQUIPMENT

Rinsing: After using TITAN Pictron EC Herbicide, empty spray unit completely and drain the whole system. Thoroughly wash inside the unit using a pressure hose. Drain the spray unit, and clean any filters in the tank, pump, lines, hoses and nozzles. After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

Decontamination: Before spraying cotton and other sensitive crops with equipment that has been used to apply TITAN Pictron EC Herbicide, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section. Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (eg. liquid Surf*, OMO*, Omomatic*, Drive* at 500mL/100L of water or the powder equivalent at 500g/100L of water) and circulate throughout the system for at least 15 minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.

RESISTANT WEEDS WARNING

TITAN Pictron EC Herbicide is a member of the pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, TITAN Pictron EC Herbicide is a Group 4 Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group 4 Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds.

GROUP **4** HERBICIDE



Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or TITAN AG Pty Ltd.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Crops susceptible to TITAN Pictron EC Herbicide include, but are not limited to: peas, lupins, lucerne, navy beans, peanuts, soybeans and other legumes, cotton, flowers, fruit, hops, ornamentals, shade trees and *Pinus* spp., potatoes, safflower, sugar beet, sunflowers, tobacco, tomatoes, vegetables and vines. TITAN Pictron EC Herbicide is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.

Picloram, one of the active constituents in this product, can remain in the soil for extended periods depending on soil type and application rate, rainfall, temperature, humidity, soil moisture and soil organic matter.

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray drift onto nearby susceptible plants/crops, cropping lands, pastures, waterways or native vegetation.

DO NOT allow physical spray drift onto waterways, native vegetation and susceptible crops.

DO NOT apply close to, or in areas, containing roots of desirable vegetation, where treated soil may be washed onto areas growing (or areas to be planted with) desirable plants.

DO NOT apply on sites where surface water from heavy rain can be expected to run off to areas containing, or to be planted with susceptible crops or plants. DO NOT move soil, which may have been treated, to areas where desirable plants are to be grown.

PROTECTION OF LIVESTOCK

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

Many plants remain poisonous after death, and stock should not be allowed to access these, as there is a likelihood that they may graze the dead material. Such material should be burnt if possible.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or watercourses with chemical or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seed. Triple rinse containers before disposal. Add rinsings to the spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (See SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (See STORAGE AND DISPOSAL section). If necessary, wash the spill area with an alkali detergent and water and absorb the wash liquid for disposal, as above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When preparing the spray, wear cotton overalls buttoned to the neck and wrists, a washable hat, elbow-length chemical resistant gloves and face shield or goggles. If the product is in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Pictron EC Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Combustible liquid. Harmful if swallowed. Causes skin irritation. Very toxic to aquatic life with long lasting effects. **Precautionary Statements:** Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking. Wash thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection/hearing protection. IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of water. Specific treatment (see on this label). If skin irritation occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. In case of fire: Use CO₂, powder or water spray to extinguish. Collect spillage. Store in a well-ventilated place. Dispose of contents/container in accordance with local/regional/national regulations.

* Other trademarks

