

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

Propyzamide 900 WG

Herbicide

ACTIVE CONSTITUENT: 900g/kg PROPYZAMIDE

GROUP **D** HERBICIDE

For selective control of certain grasses and broad leaf weeds in canola, grain legumes, legume pasture, lettuce and turf according to the Directions for use Table.

IMPORTANT: READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

APVMA Approval No.: 86871/116861

Contents: 500g-20kg



TITAN AG Pty Ltd

15/16 Princes Street, Newport NSW 2106

Tel 02 9999 6655 Fax 02 9999 0483

www.titanag.com.au

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL.

DIRECTIONS FOR USE:**Restrains:**

DO NOT apply more than once per season.

DO NOT apply to water logged soils.

DO NOT apply to land where runoff to aquatic areas or sensitive crops like cereals is likely.

CROPS	WEEDS CONTROLLED	STATE	RATE kg/ha	CRITICAL COMMENTS	
Canola	Annual Ryegrass, Winter Grass, Barley Grass, Silver Grass, Prairie Grass, Canary Grass, Great Brome, Squirrel Tail Fescue, Rats Tail Fescue	ALL STATES	0.55	Apply to moist soil prior to seeding only. Incorporate by sowing (IBS) with knife or blade points. A rainfall event (25 mm) between application and sowing, or soon after sowing is required for best results. Avoid seeding practices that moves treated soil back into the furrow.	
Grain Legumes (including lupins, chickpeas, lentils, field peas, faba beans, navy beans)				Heavy rainfall events that wash product back into the furrow can delay emergence and cause crop injury. DO NOT use on white sandy soils, or paddocks that have been recently mouldboard ploughed or spaded.	
Legume Pastures (including lucerne, clover, medics etc. grown for forage, hay or seed production)	Wild Oats, Perennial Ryegrass, Bent Grass, English Couch, Yorkshire Fog Grass, Chickweed, seedling Sorrel, Blackberry Nightshade, Patterson's Curse, seedling Wireweed	NSW, ACT, TAS, SA, WA only	0.56-0.83	Apply when weeds are at post or pre-emergent stages. SEEDLING CROPS: Apply after the third trifoliate leaf is fully developed. ESTABLISHED CROPS: Apply any time before the end of July.	
			0.83-1.11	Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended. BEST RESULTS: When applied during cool and moist conditions. Rain or irrigation is necessary to thoroughly wet the top 3 to 4cm of soil. This is similar to 25mm rain. Use rates towards the higher end of the range on heavy soils, if conditions are not optimal or where a heavy grass population is expected. Grazing or cutting PRIOR to treatment is recommended.	
Lettuce (direct sown)	Barnyard Grass, Blackberry Nightshade, Chickweed, Portulaca, Winter Grass, Rye Grass, Summer Grass, Wireweed, Prince of Wales Feather, Shepherd's Purse, Nettles	ALL STATES	2.50kg in 330-550L of water	Apply pre-emergent, immediately after sowing. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
Lettuce (transplanted)				Apply immediately after transplanting. For best results, prepare the seed bed free from clods. Sufficient irrigation to wet the top 3 to 4cm of soil should be applied within 24 hours.	
				VIC only	1.28kg in 300-550L of water
Turf and lawns Common couch	Winter Grass	ALL STATES	0.67kg (7g/100m ²)	Rate/Bowling Green 37m x 37m 94g	Apply during the autumn and winter, if established turf is infested with seedling winter grass. Ensure a uniform coverage. Irrigate immediately after application with 12-25mm irrigation or rain and again 24 hours later. Repeat treatment if necessary on Qld blue couch after 4 weeks. DO NOT RESEED FOR 60 DAYS AFTER TREATMENT. Use ONLY on established turf. DO NOT use on bluegrass, ryegrass, fescue or bent grass lawns.
Queensland blue couch		QLD only	0.33kg (3g/100m ²)	47g	
Buffalo grass		WA only	0.67kg (7g/100m ²)	94g	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**WITHHOLDING PERIODS:****CANOLA:****HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.****GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.****GRAIN LEGUMES:****HARVEST – NOT REQUIRED WHEN USED AS DIRECTED.****GRAZING – DO NOT GRAZE OR CUT FOR STOCK FOOD 12 WEEKS AFTER APPLICATION.****ALL OTHER CROPS:****DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD OR FOR SEED FOR 25 DAYS AFTER APPLICATION.****DO NOT GRAZE TREATED TURF OR LAWN; OR FEED TURF OR LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK.**

GENERAL INSTRUCTIONS

Best results are obtained by applying the product to cool, wet soil.

SPRAY PREPARATION

TITAN Propyzamide 900 WG Herbicide is a Wettable Granule formulation to be blended with water and applied as a spray. Granule formulations are designed to be added to large volumes of water and dispersed slowly under agitation. When using larger boomsprays and unable to add granules directly to a large volume in the tank via the top of the boomspray, pre-mixing may be necessary as follows:

- Partly (30%) fill the boomspray tank with water.
- Slowly add required amount of Propyzamide 900 WG Herbicide for the area to be sprayed into Granni Pot* or other pre-mixing vessel while agitating well.
- Ensure high water volumes and DO NOT add to the spray tank until the pre-mix is fully dissolved. This may take several mixes to ensure the correct amount is added to the boomspray tank.
- Add the pre-mixed TITAN Propyzamide 900 WG Herbicide into the boomspray tank with the agitation system operating.
- Fill up to the correct volume with water.
- The mixture must be kept in suspension at all times by continuous agitation.
- Tank-mixtures should be tested for physical compatibility and efficacy before use.

DO NOT mix with MCPA. If tank mixing, TITAN Propyzamide 900 WG Herbicide must be in suspension before adding other herbicides, insecticides or surfactants. Wash sprayer thoroughly after use.

DO NOT handle the water-soluble packaging with wet hands or wet gloves. Keep dry.

Equipment

Always ensure the sprayer is clean, using a suitable spray and boom cleaner as directed. It is recommended that for turf situations, up to 100L/ha spray volume is required.

RESISTANCE WEED WARNING

TITAN Propyzamide 900 WG Herbicide is a member of the amide group of herbicides with diverse sites of action. For weed resistance management this is a Group D herbicide. Some naturally occurring weed biotypes resistant to TITAN Propyzamide 900 WG Herbicide and other amide herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by TITAN Propyzamide 900 WG Herbicide or other amide herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Propyzamide 900 WG Herbicide to control the resistant weeds.

DO NOT rely exclusively on Propyzamide for weed control. Use as part of an integrated weed management program involving other modes of action and non-chemical methods of control. Advice as to strategies and alternative treatments that can be used, should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a TITAN AG representative.

RE-ENTRY PERIOD

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves.

Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply to, or allow spray to drift onto crops, pastures or non-target desirable plants, nor onto agricultural land scheduled for cropping or pasture, as injury to the crop or pasture may occur.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers or waterways by cleaning of equipment, disposal of wastes or used containers.

STORAGE AND DISPOSAL

Store in the closed original container in a dry, cool, well-ventilated area, out of direct sunlight. Keep away from contact with fertilizers, insecticides, fungicides and seeds. Protect from frost.

Single-rinse or shake remainder into spray tank. DO NOT dispose of undiluted chemicals on-site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Harmful if inhaled. Will irritate the eyes. Avoid contact with eyes and skin. DO NOT inhale dust. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves, face shield or goggles and disposable dust face mask covering mouth and nose. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Propyzamide 900 WG Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Suspected of causing cancer. Very toxic to aquatic life with long lasting effects. Precautionary Statements: Wear protective gloves/protective clothing/eye protection/face protection. Avoid release to the environment. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. IF exposed or concerned: Get medical advice/attention. Collect spillage. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations.

