

CAUTION

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

Tebuconazole 750 WDG

Fungicide

ACTIVE CONSTITUENT: 750 g/kg TEBUCONAZOLE

GROUP 3 FUNGICIDE

For the control of various diseases of bananas, peanuts, grapes, cereal crops, vegetables and other crops as specified in the Directions for Use Table.

IMPORTANT: READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

APVMA Approval No.: 64271/56286

Contents: 1kg, 5 kg, 5.71kg, 10kg, 15 kg



TITAN AG Pty Ltd

15/16 Princes Street, Newport NSW 2106

Tel 02 9999 6655 Fax 02 9999 0483

www.titanag.com.au

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL.

DIRECTIONS FOR USE

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Bananas (QLD, NSW, WA, NT only)	Leaf Spot (Yellow Sigatoka), Leaf Speckle, Black Sigatoka	130g/ha Add 3-6L water miscible oil/ha	H 1 day	<p>Maintain good deleafing practices to reduce disease inoculum. Very old leaves, and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of TITAN Tebuconazole 750 WDG Fungicide.</p> <p>Tropical areas (e.g. North Qld, NT, Ord River): Apply a regular schedule of protectant sprays. When conditions favour disease, apply a maximum of 2 consecutive TITAN Tebuconazole 750 WDG Fungicide sprays at 14 day intervals. DO NOT apply more than 6 TITAN Tebuconazole 750 WDG Fungicide sprays in any 12 month period. DO NOT apply any TITAN Tebuconazole 750 WDG Fungicide sprays in the months of July, August, September and October.</p> <p>Sub-tropical areas (e.g. South Qld, NSW): Commence spraying with TITAN Tebuconazole 750 WDG Fungicide at the onset of warm and humid/wet weather, normally December. Repeat at 21 to 28 day intervals using a maximum of 2 consecutive TITAN Tebuconazole 750 WDG Fungicide sprays. DO NOT apply more than 5 TITAN Tebuconazole 750 WDG Fungicide sprays in any 12 month period. TITAN Tebuconazole 750 WDG Fungicide is approved for use in banana plantations interplanted with avocados.</p> <p>Ground application: Apply by misting machine or airblast sprayer in a convenient volume of water.</p>
Grapevines	Powdery Mildew	<p>Dilute Spraying 17.2g/100L</p> <p>Concentrate spraying Refer to 'Special instructions for grapevines' in the Application section</p>	H 8 weeks	<p>Apply thoroughly as part of the following 5 spray program:</p> <ol style="list-style-type: none"> 1. when shoots 10-20 cm long. 2. pre-flowering. 3. flowering. 4. after fruit set. 5. before bunches close. <p>This use is subject to a CropLife Australia Fungicide Resistance Management Strategy. DO NOT apply more than two consecutive sprays of TITAN Tebuconazole 750 WDG Fungicide. DO NOT apply more than three sprays of TITAN Tebuconazole 750 WDG Fungicide per season. DO NOT use TITAN Tebuconazole 750 WDG Fungicide curatively.</p> <p>Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use in equipment that requires concentrate rates greater than 150mL/100L of water (5x). DO NOT apply in volumes less than 250 /ha.</p> <p>Add a non-ionic wetting agent at 10mL/100L regardless of whether applying by dilute or concentrate spraying.</p>
Green beans	Rust	200g/ha + Agridex* 1L/ha	G 3 days H 3 days	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high. Applied by ground rig only.
Lettuce	Sclerotinia Rot	200g/ha	H 5 weeks	<p>Apply by boom spray. Apply only during the early stages of plant development (note the 5 week WHP). Apply a maximum of 2 applications at 7-10 day intervals in rotation with other chemicals registered for this use.</p> <p>Apply only to field crops. DO NOT apply in greenhouse (i.e. protected cropping) situations or hydroponically grown lettuce crops. Control of Sclerotinia in lettuce should not be solely reliant on fungicides. Other control measures such as crop rotations, resistant varieties and planting techniques should be employed.</p>
Onions (TAS only)	White Root Rot	1g/100 m of row mixed with 145-218g lime super/100 m of row	—	<p>Before sowing, apply TITAN Tebuconazole 750 WDG Fungicide onto lime super. Ensure good coverage of all lime super particles. Apply TITAN Tebuconazole 750 WDG Fungicide treated lime super when sowing onion seed. Seed and lime super can either be mixed in the same box on the drill or placed in different boxes and sown down the same tube. Apply in a band width of 2cm. Ensure that the correct rate of TITAN Tebuconazole 750 WDG Fungicide is used otherwise some delay in emergence and reduced stands of seedlings may occur.</p>

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Pawpaw	Black Spot	170g/ha	H 3 days	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Spray equipment must be properly calibrated to apply the correct amount of TITAN Tebuconazole 750 WDG Fungicide. Apply TITAN Tebuconazole 750 WDG Fungicide at 14 day intervals. Ensure thorough coverage of leaves and fruit. Alternate TITAN Tebuconazole 750 WDG Fungicide with sprays of a protectant fungicide (e.g. Dithane DF). DO NOT apply more than 6 sprays of TITAN Tebuconazole 750 WDG Fungicide (or any Group 3 fungicide) on any block in any 12 month period.
Peanuts (South QLD, NSW only)	Early Leaf Spot, Late Leaf Spot, Rust, Net Blotch	Low disease 100g/ha + Agridex 1L/ha	H 3 weeks G 3 weeks	Regularly check high risk areas in the crop for disease (eg lower leaves, shaded plants). When leaf spot or net blotch or rust can easily be found then either; – spray immediately after the last disease conducive weather (eg rain or heavy dews) OR – spray before the next disease conducive weather. Repeat after 14 days if conditions remain favourable to disease development. If not, repeat before or just after the next disease conducive weather. Ground application: Apply in at least 100L of water/ha. Use of TITAN Tebuconazole 750 WDG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. TITAN Tebuconazole 750 WDG Fungicide) before changing to a fungicide of a different activity group. Apply a maximum of 5 Group 3 sprays per season.
		High disease OR wet weather 166g/ha + Agridex 1L/ha		
		High disease AND wet weather 250g/ha + Agridex 1L/ha		
Peanuts (North Qld, WA, NT only)	Early Leaf Spot, Late Leaf Spot, Rust	Low disease 132g/ha + Agridex 1L/ha	H 21 days G 21 days	Begin spraying at 3-4 weeks after planting. If band spraying, apply half the appropriate rate as a 45 cm band directly over the row. Sprays should not be banded after 6 weeks from planting. Repeat applications at 14 day intervals. If prolonged wet weather or heavy rains occur, shorten spray interval to 10-12 days. Ground application: Apply in at least 100L of water/ha. Use of TITAN Tebuconazole 750 WDG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. TITAN Tebuconazole 750 WDG Fungicide) before changing to a fungicide of a different activity group. Apply a maximum of 5 Group 3 sprays per season.
		Moderate disease 166g/ha + Agridex 1L/ha		
		Severe disease 250 g/ha + Agridex 1L/ha		
	Net Blotch	Low disease 166 g/ha + Agridex 1L/ha		Apply at 14 day intervals. If prolonged cool moist weather occurs, shorten spray interval to 10-12 days. Ground application: Apply in at least 100L of water/ha. Use of TITAN Tebuconazole 750 WDG Fungicide on peanuts is subject to a CropLife Australia Fungicide Resistance Management Strategy. Apply no more than 3 consecutive Group 3 sprays alone (eg. TITAN Tebuconazole 750 WDG Fungicide) before changing to a fungicide of a different activity group. Apply a maximum of 5 Group 3 sprays per season.
		Moderate - severe disease 250g/ha + Agridex 1L/ha		
Peas	Powdery Mildew	83g/ha	H 3 days G 3 days	Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under some conditions. Ground application: Apply in at least 50L of water/ha.
Pyrethrum	<i>Sclerotinia sclerotiorum</i>	200g/ha	–	Apply twice, in rotation with other control measures, at 7-10 day intervals. Commence at 1 to 2% flowering. Use under direction of pyrethrum advisers.
Ryegrass and fescue seed crops	Leaf Rust, Stem Rust	166g/ha	–	Monitor crops closely and spray at the first signs of disease. Continuing disease pressure or reinfection may require a further application 3-4 weeks later. Apply in at least 100L of water/ha. Ensure thorough coverage, and use higher water volumes in dense or advanced crops.

CROP	DISEASE	RATE	WHP	CRITICAL COMMENTS
Sugar cane (variety Q124 only)	Orange Rust	166g/ha + Agridex 1L/ha	H 4 weeks G 4 weeks	Even low levels of orange rust suppress yields so it is important to apply TITAN Tebuconazole 750 WDG Fungicide early in the development of the disease epidemic. Begin monitoring disease levels early. Check crops at least weekly when climatic conditions favour the development of disease. Apply as a foliar spray when disease begins to escalate rapidly. Repeat application after 14 days if conditions remain favourable to orange rust spore germination. DO NOT apply more than 2 TITAN Tebuconazole 750 WDG Fungicide sprays per season. DO NOT apply if heavy rains or storms that are likely to cause surface runoff are forecast with greater than 50% probability within 24 hours (48 hours if possible) of application. Ground application: Use droppers and directed sprays and sufficient water volume to ensure thorough coverage.
Wheat	Leaf Rust, Stripe Rust, Septoria Nodorum Blotch, Yellow Leaf Spot	83 or 166g/ha	H 5 weeks G 14 days	Use higher rate when longer disease control is required. Stripe Rust: See spray timings under General Instructions. Other diseases: Apply from full flag leaf emergence to early head emergence. The addition of mineral crop oil (e.g. D-C-Trate or equivalent) at 1%, may improve performance of TITAN Tebuconazole 750 WDG Fungicide on wheat, oats and barley. Ground application: Apply in at least 50L of water/ha.
	Septoria Tritici Blotch	166 g/ha		
Oats	Crown Rust	83 or 166g/ha		
Barley	Scald	83g/ha	H 5 weeks G 14 days	Apply at late tillering to early jointing. Ground application: Apply in at least 50L of water/ha.
	Powdery Mildew	83 or 166 g/ha		Apply when 5% of the leaf area is infected. Use higher rate when longer disease control is required. Ground Application: Apply in at least 50L of water/ha.
Wheat Oats	Stem Rust		H 5 weeks G 14 days	Apply if more than 5% of stems become infected between full flag leaf emergence to late flowering. Where stem rust is the major disease, yield responses are usually optimised by delaying application until full head emergence, and using the higher rate. In severe cases, if a majority of stems are infected prior to full head emergence, apply at 83 g/ha as soon as possible and if necessary, repeat after 3 weeks when heads are fully emerged. Ground application: Apply in at least 50L of water/ha.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS H = harvest, G = grazing

Bananas, Avocados: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.

Beans, Peas: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 3 DAYS AFTER APPLICATION.

Cereals: DO NOT HARVEST FOR 5 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Grapevines: DO NOT HARVEST FOR 8 WEEKS AFTER APPLICATION.

Onions: NOT REQUIRED WHEN USED AS DIRECTED.

Pawpaw: DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

Peanuts: DO NOT HARVEST FOR 21 DAYS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

Sugar cane: DO NOT HARVEST FOR 4 WEEKS AFTER APPLICATION.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Special warning – bananas

There are certain conditions when the surface of recently emerged fruit is particularly prone to marking damage from spray applications. In circumstances where application will be made to very rapidly growing fruit in hot conditions with strong direct light, it is recommended that all emerged bunches be bagged prior to spraying to minimise risk of fruit marking. Never include adjuvants other than water miscible oils with Tebuconazole sprays. The inclusion of wetting agents is known to cause phytotoxicity to young fruit.

Foliar diseases on cereal crops

DO NOT apply to cereal crops more than once per season (166g/ha rate) or more than twice per season (83g/ha rate). Treatment will give approximately three weeks disease suppression. Economic responses may not be gained by spraying crops past flowering stage. The effects of fungicide application will not be clearly seen for 7-10 days after application.

Spray timings for Stripe Rust control

Obtain advisory literature from Department of Agriculture for classification of resistant and susceptible varieties. The following spray program is suggested:

Seedling infections:

When approximately 20 out of 100 leaves show first signs of infection during tillering to jointing, apply a spray within one week.

Adult infections - susceptible varieties:

When approximately 10 out of 100 leaves show first sign of infection, apply a spray within one week. DO NOT delay.

Adult infections - moderately susceptible varieties:

When approximately 15 to 20 leaves out of 100 leaves show first sign of infection, apply a spray within one week. DO NOT delay.

Adult infections - moderately resistant and resistant varieties:

Monitor carefully. If rust appears and spreads, spray within one week.

FUNGICIDE RESISTANCE WARNING

TITAN Tebuconazole 750 WDG Fungicide is a member of the DMI group of fungicides. For fungicide resistance management the product is a Group 3 fungicide. Some naturally occurring individual fungi resistant to the product and other Group 3 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product and other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that result from failure of this product to control resistant fungi.



EXPORT OF TREATED PRODUCE

Growers should note that MRLs or import tolerances do not exist in all markets for edible produce treated with TITAN Tebuconazole 750 WDG Fungicide. If you are growing edible produce for export, please check with TITAN AG Pty Ltd for the latest information on MRLs and import tolerances before using TITAN Tebuconazole 750 WDG Fungicide.

MIXING

Shake container vigorously, then add the required quantity of TITAN Tebuconazole 750 WDG Fungicide to water in the spray vat while stirring or with agitators in motion. Add the required amount of Agridex* (beans, peanuts, sugar cane) or water miscible oil (bananas) and mix thoroughly.

SPECIAL INSTRUCTIONS FOR GRAPEVINES

Dilute Spraying

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of runoff. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate Spraying

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY

1. Dilute spray volume as determined above: For example 1500L/ha
2. Your chosen concentrate spray volume: For example 500L/ha
3. The concentration factor in this example is:
 $3 \times (\text{i.e. } 1500\text{L} \div 500\text{L} = 3)$
4. As the dilute label rate is 30mL/100L, then the concentrate rate becomes
 3×30 , that is 90mL/100L of concentrate spray.

- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- DO NOT use a concentrate rate higher than that specified in the Critical Comments.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

PRECAUTIONS

Re-entry Period: DO NOT enter treated area until spray has until spray has dried after treatment unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each days use.

DO NOT enter treated areas for high exposure activities such as grape girdling for nine days after application unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, drains or waterways with the chemical or used containers. A spraydrift minimisation strategy should be employed at all times. Spray drift may occur under adverse meteorological conditions or from certain spraying equipment. DO NOT allow spray to drift onto sensitive areas including, but not limited to, susceptible plants/crops, cropping land, pasture, natural streams, rivers, wetlands, waterways or human dwellings.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management authority is not available, bury the empty packaging 500mm the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container, preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist or equivalent clothing and a washable hat and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Tebuconazole 750 WDG Fungicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if swallowed. Suspected of damaging fertility or the unborn child. Toxic to aquatic life with long lasting effects. **Precautionary Statements:** Wear protective gloves/protective clothing/eye protection/face protection. Avoid release to the environment. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. IF exposed or concerned: Get medical advice/attention. Rinse mouth. Collect spillage. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations.

