

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

TERBUTRYN 500 SC

HERBICIDE

ACTIVE CONSTITUENT: 500g/L TERBUTRYN

GROUP **5** HERBICIDE



Controls certain early competing broadleaf weeds in wheat, barley, triticale, pastures, some varieties of oats and field peas as per Directions for Use table.

APVMA Approval No.: 67767/56821

Pack Size: 1L; 5L-1000L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | Fax (02) 9999 0483
titanag.com.au

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL

DIRECTIONS FOR USE

Restrains: DO NOT APPLY THIS PRODUCT WHEN TEMPERATURES ARE GREATER THAN 18°C FOR THE DAY PRIOR TO, THE DAY OF AND FOR THE 4 DAYS FOLLOWING APPLICATION.

DO NOT spray under dry frosty conditions or if very dry conditions prevail.

DO NOT spray if weeds or crop/pasture are affected by insects, disease, dust or adverse weather conditions.

DO NOT add wetting agents or crop oils.

DO NOT use main or nozzle filters finer than 50 mesh for tank mixes.

DO NOT tank mix with selective grass herbicides.

DO NOT apply tank mixture of TERBUTRYN plus triasulfuron (750g/kg) WG if another Group B Herbicide (ASL inhibitor) has been used during the current season.

IF RAIN FALLS within 6 hours of application, the effect could be diminished.

CROP	WEEDS CONTROLLED	STATE	RATE PER HA	CRITICAL COMMENTS
Wheat, Barley, Triticale and some varieties of Oats: from 3 leaf to early tillering stage	Amsinckia (Yellow Burrweed), Capeweed, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Fumitory (Common, White and Dense Flowered), Paterson's Curse, Toad Rush – 2 to 6 leaves	NSW, VIC only	550 to 850mL	Use the highest rate when Deadnettle or Toad Rush is the major problem. Apply to boom spray or aircraft.
	The above weeds in addition to Charlock, Doublegee (Three-cornered Jack), Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Turnip Weed, Wild Turnip – 2 to 6 leaves	NSW, VIC only	550 to 850mL + 300mL of a 500g/L 2,4-D (as amine) or 300mL of a 500g/L MCPA as a tank mix	DO NOT add MCPA or 2,4-D if undersown Medics are present. Apply by boom spray or aircraft. Use the highest rate when Deadnettle, Doublegee (Three-cornered Jack) or Toad Rush is the major problem.
	<i>Plagiobothrys canascens</i> – 2 to 6 leaves	NSW, VIC only	850mL	
	Turnip Weed	NSW only	300mL + 6.5g of a 750g/kg WG triasulfuron	Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip		300mL + 6.5g of a 750g/kg WG triasulfuron to 440mL + 10g of a 750g/kg WG triasulfuron	
	Black Bindweed (Climbing Buckwheat), Faba Bean (volunteer), Field Pea (volunteer), Medic, Shepherd's Purse		440mL + 10g of a 750g/kg WG triasulfuron	
	Deadnettle, Mexican Poppy, Wild Radish		440mL + 10g of a 750g/kg WG triasulfuron to 600mL + 13g of a 750g/kg WG triasulfuron	
	Coreopsis, Sunflower (volunteer), Vetch		600mL + 13g of a 750g/kg WG triasulfuron	
	Paterson's Curse, Turnip Weed		300mL + 6.5g of a 750g/kg WG triasulfuron	
	Hedge Mustard, Indian Hedge Mustard, London Rocket, Wild Turnip		300mL + 6.5g of a 750g/kg WG triasulfuron to 440mL + 10g of a 750g/kg WG triasulfuron	
	Crassula, Faba Bean (volunteer), Hyssop Loosestrife, Medic, Prickly Lettuce (Whipthistle)	VIC only	440mL + 10g of a 750g/kg WG triasulfuron	
	Deadnettle, Field Pea (volunteer), Wild Radish		440mL + 10g of a 750g/kg WG triasulfuron to 600mL + 13g of a 750g/kg WG triasulfuron	
	Lupins (volunteer), Vetch, Wire Weed (less than 3 leaves) suppression		600mL + 13g of a 750g/kg WG triasulfuron	
	Turnip Weed		300mL + 6.5g of a 750g/kg WG triasulfuron	
	Dense flower Fumitory, Indian Hedge Mustard, Small flower Fumitory, Wild Turnip	QLD only	300mL + 6.5g of a 750g/kg WG triasulfuron to 440mL + 10g of a 750g/kg WG triasulfuron	Spray when weeds are in the 2 to 6 leaf stage, except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Best results are obtained when good soil moisture has been present since planting.
	Black Bindweed (Climbing Buckwheat), London Rocket		440mL + 10g of a 750g/kg WG triasulfuron	
	Wild Radish		440mL + 10g of a 750g/kg WG triasulfuron to 600mL + 13g of a 750 g/kg WG triasulfuron	
	Coreopsis, Corn Gromwell (Sheepweed or White Ironweed)		600mL + 13g of a 750g/kg WG triasulfuron	

CROP	WEEDS CONTROLLED	STATE	RATE PER HA	CRITICAL COMMENTS
Wheat, Barley, Triticale and some varieties of Oats: from 3 leaf to early tillering stage – <i>continued</i>	Amsinckia (Yellow Burrweed), Capeweed, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Doublegee (Three-cornered Jack), Fumitory (Common, White and Dense Flowered), Rough Poppy, Salvation Jane, Toad Rush – 2 to 6 leaves	SA only	550 to 850mL	DO NOT add MCPA or 2,4-D if undersown Medics are present. Apply by boom spray or aircraft. Use the highest rate when Deadnettle, Doublegee (Three-cornered Jack) or Toad Rush is the major problem. DO NOT apply to disease infected wheat as yellowing may occur. Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	The above weeds in addition to Charlock, Hedge Mustard, Hexham Scent, Indian Hedge Mustard, Mallee Catch-Fly, Turnip Weed, Wards Weed, Wild Turnip – 2 to 6 leaves		550 to 850mL + 300mL of a 500g/L 2,4-D (as amine) or 300mL of a 500g/L MCPA as a tank mix	
	Faba Beans (volunteer), Long Fruited Turnip, Long Headed Poppy, Paterson's Curse, Turnip Weed		300mL + 6.5g of a 750g/kg WG triasulfuron	
	Hedge Mustard, Indian Hedge Mustard, Wild Turnip		300mL + 6.5g of a 750g/kg WG triasulfuron to 440mL + 10g of a 750g/kg WG triasulfuron	
	Ball Mustard, Crassula, Medic, Prickly Lettuce (Whipthistle)		440mL + 10g of a 750g/kg WG triasulfuron	
	Field Pea (volunteer), Wild Radish		440mL + 10g of a 750g/kg WG triasulfuron to 600mL + 13g of a 750g/kg WG triasulfuron	
	Doublegee (Spiny Emex, Three-cornered Jack), Lupins (volunteer), Vetch, Wireweed (less than 3 leaves) suppression		600mL + 13g of a 750g/kg WG triasulfuron	
	Capeweed, Corn Gromwell, Fumitory, Yellow Burrweed – 2 to 6 true leaves, Doublegee (Three-cornered Jack), Toad Rush – 1 to 4 leaves	WA only	850mL	Use the highest rate when Fumitory is the major problem weed in addition to Wild Turnip, Indian Hedge Mustard, London Rocket, Lupin and Wild Radish. Spray when weeds are in the 2 to 6 leaf stage (up to 6 node/leaflet for Field Peas, 10 leaflet for Vetch), except 2 to 4 leaf for Doublegee. Use rates towards the lower end of the range (where applicable) when weeds are small and soil conditions are very moist. Spray only when weeds are actively growing. Spray only after good rain and when top soil is moist. Best results are obtained when good soil moisture has been present since planting.
	The above weeds in addition to Indian Hedge Mustard, London Rocket, Lupin, Wild Radish and Wild Turnip – 2 to 6 leaves		550 to 850mL or 600mL of a 500g/L MCPA as a tank mix	
	Turnip Weed		300mL + 6.5g of a 750g/kg WG triasulfuron	
	Denseflower Fumitory, Hedge Mustard, Smallflower Fumitory		300mL + 6.5g of a 750g/kg WG triasulfuron to 440mL + 10g of a 750g/kg WG triasulfuron	
	Ball Mustard, Field Pea (volunteer)		440mL + 10g of a 750g/kg WG triasulfuron	
	Deadnettle		440mL + 10g of a 750g/kg WG triasulfuron to 600mL + 13g of a 750g/kg WG triasulfuron	
	Australian Crassula, Doublegee (Spiny Emex, Three-cornered Jack), Lupins (volunteer)		600mL + 13g of a 750g/kg WG triasulfuron	
Pastures Use on mixtures of either/or Sub Clover (min. of 3 to max. of 10 leaves), Balansa Clover (established plants), White Clover (min. of 5 to max. of 10 leaves) with either/or Ryegrass, Phalaris, Cocksfoot (min. 3 leaves)	Capeweed (seedling to 10cm diameter), Paterson's Curse (seedling to 10cm diameter)	NSW, VIC, TAS only	550 to 700mL	Apply by boom spray or aircraft. Use the lowest rate early in the season or when weeds are small and young. Increase rate as weed size and age increases. Early germinating Paterson's Curse with strong root systems may not be controlled. Some temporary damage may be seen on clover if adverse growing conditions occur at end after spraying. Where a selective grass herbicide or simazine is to be or has been used, TITAN Terbutryn 500 SC Herbicide should not be used in the same season. See Pasture Safety section in General Instructions.
	Capeweed (seedling to 20cm diameter), Crassula (1 to 5cm diameter), Erodium botrys (seedlings to 8cm diameter), Paterson's Curse (seedling to 20cm diameter), Toad Rush (1 to 5 leaves)		400 to 700mL plus 300 to 500mL of MCPA 500 amine as a tank mix	

CROP	WEEDS CONTROLLED	STATE	RATE PER HA	CRITICAL COMMENTS
Field Peas	Amsinckia (Yellow Burrweed), Capeweed, Common Fumitory, Corn Gromwell (Sheep Weed or White Iron Weed), Deadnettle, Paterson's Curse, Toad Rush	VIC only	700 to 850mL	Apply as an early post-emergent application when the peas are 10 to 12cm high and have developed 3 to 5 nodes. Weeds should be young and actively growing and in the 4 to 5 true leaf stage. Use rates towards the higher end of the scale when Deadnettle is the major problem and when weeds are in the 6 true leaf stage. Where a selective grass herbicide is to be used, apply TITAN Terbutryn 500 SC Herbicide 2 weeks before the application of the selective herbicide.
Canning Peas (Green Peas, Vining Peas, Processing Peas)	Blackberry Nightshade, Capeweed, Chickweed, Common Fumitory, Corn Gromwell, Deadnettle, Fat Hen, Paterson's Curse, Toad Rush, Yellow Burrweed	TAS only	750mL to 1L	Apply as an early post-emergent application when the peas are 10 to 12cm high and have developed 3 to 6 nodes and the weeds are young and actively growing and in the 2 to 4 true leaf stage. Use rates towards the higher end of the scale when Deadnettle and Blackberry Nightshade are the major problems.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

FIELD PEAS, CANNING PEAS: DO NOT APPLY WITHIN 4 WEEKS OF HARVEST.

PEA FODDER AND FORAGE: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

PASTURES: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

**CEREALS – TERBUTRYN ALONE WITH TERBUTRYN PLUS MCPA OR 2,4-D AS A TANK MIX:
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**CEREALS – TERBUTRYN PLUS TRIASULFURON (750 g/kg WG) TANK MIX:
DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

GENERAL INSTRUCTIONS

IMPORTANT

Spray only after good rain and when top soil is moist and when weeds are actively growing. Best results are obtained when good soil moisture has been present since planting. DO NOT apply during dry or waterlogged conditions.

RESISTANCE WEEDS WARNING

TITAN Terbutryn 500 SC Herbicide is a member of the Triazine group of herbicides. This product has the photosystem II photosynthesis inhibitor mode of action. For weed resistance management TITAN Terbutryn 500 SC Herbicide is a Group 5 herbicide. Some naturally-occurring weed biotypes resistant to TITAN Terbutryn 500 SC Herbicide and other Group 5 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by TITAN Terbutryn 500 SC Herbicide or other Group 5 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Terbutryn 500 SC Herbicide to control resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a TITAN AG representative.

CROP SAFETY

DO NOT apply this product to Avon, Saia, Barmah and Cassia varieties of oats. Before applying to other varieties of oats, growers should seek advice from TITAN AG for crop safety information. Undersown Medics and American Lucerne varieties may be damaged if sprayed. With early and late sown crops damage may occur if spraying is undertaken when temperatures are greater than 18°C. Avoid spraying crops prior to May and from mid-August onwards as daily temperatures and weed size may not be satisfactory. Early sown crops should not be sprayed before the onset of frost and cooler weather.

PASTURE SAFETY

DO NOT apply this product to pasture species other than those stated in the "Directions for Use". DO NOT apply to medics and lucerne as damage may occur. DO NOT apply to irrigated pasture or seed crops as damage may occur. DO NOT apply on pastures prior to May or after mid-August or when temperatures are greater than 18°C for the day prior to, the day of and for 4 days following application as pasture damage will occur. DO NOT apply prior to or following the use of a grass selective herbicide or simazine in the same season as damage to pasture may occur. DO NOT use in situations where oversowing of pastures may occur. Pastures which contain young species should be sprayed using the lower label rates. Increase product rate as pasture age and size increases. Only apply this product when conditions at and following application are favourable to pasture growth.

MIXING

Stir product or invert container several times before opening as settling of contents may occur after storage for some weeks. Pour the product into the spray vat through a strainer to remove any dry particles or flakes, which can occasionally occur under hot storage conditions. Add the full quantity of product to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part used containers immediately. If tank mixing, add TITAN Terbutryn 500 SC Herbicide to the spray tank first, then other products. Maintain constant agitation and use immediately.

APPLICATION

Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is obtained. Aim to apply 50 to 110L per hectare.

Aircraft Application: Always spray in a cross wind of less than 5 knots, or use smoke generator to indicate air movement. Ensure good spray coverage is obtained. For maximum coverage use fine nozzles. Apply 20 to 30L per hectare.

Crop Rotation Guidelines

– **TERBUTRYN plus triasulfuron (750g/kg) WG tank mixture**

The following crops can be planted after:

In areas where pH is less than 7.5

– 9 months after an application; barley, cereal rye, chickpeas, faba beans, field peas, lupins, medics, oats, canola, safflower, subclover, triticale, wheat.

In areas where pH is greater than 7.5

– 9 months after an application; barley, cereal rye, oats, triticale, wheat.

– 14 months after an application: cotton, maize, sorghum, soybeans, sunflowers. Where residual herbicides are applied following an application of TERBUTRYN plus Triasulfuron (750g/kg) WG, the crop rotational guidelines for these products must be followed.

– 22 months after an application: chickpeas, faba beans, field peas, lupins, medics, canola, safflower, subclover.

For all other crops advice should be sought from AGROREG.

COMPATIBILITY

This product is compatible with Ally*, MCPA 500 Amine (not LVE), 2,4-D Amine, Triasulfuron (750 g/kg) WG, Banvel* 200, Banvel* M, Cadence* and Supracide*.

PRECAUTION

Re-entry Period: DO NOT enter treated area until spray has dried.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate ponds, waterways or drains with the chemical or used container.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT use in situations where off-target damage may occur, eg vines, tree crops. If using 2,4-D amine or MCPA apply with equipment which will not be used in hormone susceptible crops, particularly cotton and vines. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

For Refillable containers: Empty contents fully into application equipment. Close all vales and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Avoid contact with skin. Repeated exposure may cause allergic disorders. When using the product wear elbow-length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Terbutryn 500 SC Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Toxic to aquatic life with long lasting effects. Precautionary Statements: Avoid release to the environment. Collect spillage. Dispose of contents/ container in accordance with local/regional/national regulations.

