

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

IMAZAPIC 240

HERBICIDE

ACTIVE CONSTITUENT: 240g/L IMAZAPIC present as the AMMONIUM SALT

GROUP **2** HERBICIDE

For the pre-emergence control of certain annual grass and broadleaf weeds in fallow situations, Sugarcane and Peanuts and early post-emergence control of certain annual grass and broadleaf weed in Peanuts and Sugarcane as specified in the Directions For Use table.

APVMA Approval No.: 82379/105606

Pack Size: 5L-20L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | titanag.com.au

SL	Formulation Type Soluble Liquid Concentrate
IN A TRANSPORT EMERGENCY • DIAL 000 • POLICE OR FIRE BRIGADE	
TRANSPORT AND HANDLING NOT A DANGEROUS GOOD ACCORDING TO THE AUSTRALIAN DANGEROUS GOODS (ADG) CODE FOR TRANSPORT BY ROAD AND RAIL	

DIRECTIONS FOR USE

Restraints:

DO NOT apply to soils saturated with water.

DO NOT apply by aircraft.

SITUATION	WEEDS CONTROLLED	RATE/ha	CRITICAL COMMENTS
Fallow prior to planting winter crops as directed in the follow crops table (QLD and Nthn NSW only)	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Stink Grass (<i>Eragrostis cilianensis</i>), Blowaway Grass (<i>Panicum decompositum</i>), Button Grass (<i>Dactyloctenium radulans</i>), Pigweed (<i>Portulaca oleracea</i>), Yellow Vine (<i>Tribulus terrestris</i>), Mintweed (<i>Salvia reflexa</i>), Boggabri Weed (<i>Amaranthus mitchelli</i>), Dwarf Amaranth (<i>Amaranthus macrocarpus</i>), Peachvine/Cowvine (<i>Ipomoea lonchophylla</i>)	150 to 200mL	Treatments may be applied from the end of the previous crop harvest but not less than 3 months prior to planting of winter crop (refer to follow crops table). Best results are obtained where treatments are applied to dry, weed-free soil prior to weed germination. Treatments can be applied to standing crop residues. Sufficient rainfall is required after application and prior to weed emergence to wet soil to a depth of 5cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Emerged weeds must be controlled by prior cultivation, or by application of a knockdown herbicide. Where infestation of emerged weeds is light, tank mix treatment with glyphosate or paraquat. Cultivation following application of treatments may reduce pre-emergence weed control provided by TITAN Imazapic 240 Herbicide. Prolonged wet soil conditions following treatment will reduce the effective life of pre-emergence treatments. Weed escapes may require follow up application of knockdown herbicides. DO NOT apply more than 200mL/ha per season on cropland. DO NOT apply in more than two successive seasons on cropland. DO NOT overspray headlands.
Sugarcane (QLD, NSW, WA only)	Awnless Barnyard Grass (<i>Echinochloa colona</i>), Barnyard Grass (<i>E. crus-galli</i>), Blackberry Nightshade (<i>Solanum nigrum</i>), Blue Billygoat Weed (<i>Ageratum houstonianum</i>), Common Sida (<i>Sida rhombifolia</i>), Giant/Black Pigweed (<i>Trianthema portulacastrum</i>), Glossy Nightshade (<i>Solanum americanum</i>), Green Amaranth (<i>Amaranthus viridis</i>), Green Summer Grass (<i>Brachiaria subquadrifida</i>), <i>Ipomoea</i> spp., Guinea Grass (<i>Panicum maximum</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Milkweed (Mexican Fire Plant) (<i>Euphorbia heterophylla</i>), Pigweed (<i>Portulaca oleracea</i>), Summer Grass (<i>Digitaria ciliaris</i>)	300 to 400mL	DO NOT apply more than once a year to the same crop. Pre-emergence application: PLANT CANE: Apply after planting and before emergence of the cane. On sandy soil, ensure thorough coverage of setts to a minimum depth of 5cm. RATOON: Apply after harvest and before new cane growth appears. Use the higher rates of TITAN Imazapic 240 Herbicide for increased residual control of weeds or anticipated high weed pressure. Heavy trash layers may prevent the product from reaching the soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5cm. Emerged weeds must be controlled by prior cultivation, or application of an appropriate herbicide. Weed control can be improved under dry conditions or where incorporating moisture has been insufficient, by cultivating the treated soil above setts. Care needs to be taken to minimise exposure of untreated soil or setts. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions. Heavy rain and/or irrigation within 2 days following application may concentrate TITAN Imazapic 240 Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Weed escapes may require follow up application of an appropriate herbicide. Contact TITAN AG Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label).
	As above plus Crowsfoot Grass (<i>Eleusine indica</i>)	400mL plus 2.3L Farmalinx Panda 435	Incorporate into the soil within 7 days of application, either by overhead irrigation, rainfall or mechanical incorporation, to a minimum depth of 5cm. Read all directions for use in sugarcane as well as the Restraints and Follow Crops restrictions on the Farmalinx PANDA 435 label.
	As above plus control of those weeds expected with these rates of 250g/L paraquat products alone.	300 to 400mL plus 1.2 to 1.6L of 250g/L paraquat product	Post-emergence application (in emerged cane): ALWAYS tank mix TITAN Imazapic 240 Herbicide with registered and appropriate rates of paraquat for the emerged weeds. Use the higher rates of TITAN Imazapic 240 Herbicide for increased residual control of weeds or anticipated high weed pressure. Use the higher rates of paraquat for dense, more mature weeds. Apply as a carefully directed inter-row spray to weeds and at the base of the plant, minimising contact with sugarcane foliage. Contact with Sugarcane foliage may cause temporary crop damage observed as interveinal yellowing and reduction in crop biomass or crop height. Refer to the Paraquat label re spray boom types and precautions. Reduce the incidence of "fines" by using the correct nozzles and operating pressures. Replace worn or old nozzles and calibrate equipment on a regular basis. DO NOT cultivate within 1 hour of application. Heavy rain and/or irrigation within 2 days following application may concentrate TITAN Imazapic 240 Herbicide in the furrow and cause temporary yellowing and stunting of cane leaves. DO NOT spray plants covered with heavy dew. Prolonged wet soil and/or cool conditions can increase crop damage, especially in sandy soil. Normal growth will return when conditions become warm and drier. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions. Contact TITAN AG Pty Ltd for advice on variety susceptibility before using this product (refer to the CROP SAFETY section of the label).

SITUATION	WEEDS CONTROLLED	RATE/ha	CRITICAL COMMENTS
Peanuts (QLD, NSW, NT only)	Awnless Barnyard Grass <i>(Echinochloa colona)</i> , Barnyard Grass (<i>E. crus-galli</i>), Blackberry Nightshade <i>(Solanum nigrum)</i> , Blue Billygoat Weed <i>(Ageratum houstonianum)</i> , Common Sida (<i>Sida rhombifolia</i>), Giant/Black Pigweed <i>(Trianthema portulacastrum)</i> , Glossy Nightshade <i>(Solanum mericanum)</i> , Green Amaranth (<i>Amaranthus viridis</i>), Green Summer Grass <i>(Brachiaria subquadrifera)</i> , <i>Ipomoea</i> spp., Guinea Grass (<i>Panicum maximum</i>), Liverseed Grass (<i>Urochloa panicoides</i>), Milkweed (Mexican Fire Plant) <i>(Euphorbia heterophylla)</i> , Pigweed (<i>Portulaca oleracea</i>), Summer Grass (<i>Digitaria ciliaris</i>)	300 to 400mL	Pre-emergence application: Optimum weed control will occur when moisture is maintained in the top 5cm of soil. Apply to weed-free soil prior to weed germination. Sufficient rainfall, or overhead irrigation, is required after application and prior to weed emergence to wet soil to a minimum depth of 5cm. Use the higher rates for increased residual control of weeds or anticipated high weed pressure. Pre-emergence weed control may be limited on Krasnozem or red brown Ferrosol soil types, where moisture is not maintained in the top 5cm of soil. Refer to Soil Type in the General Instructions. DO NOT apply to any one crop more than once per season.
	As above plus Nutgrass* <i>(Cyperus rotundus)</i>	400mL plus Hasten* or Kwicken* at 1L/100L (or a similar esterified oil adjuvant)	Post-emergence application: Apply prior to crop flowering. DO NOT apply to any one crop more than once per season. * Nutgrass: Suppression only when applied to nutgrass at the 3 to 4 leaf stage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

HARVEST: NOT REQUIRED WHEN USED AS DIRECTED.

GRAZING/STOCKFEED WHP:

Wheat, Barley, Chickpea, Peanuts: DO NOT GRAZE TREATED VEGETATION, STUBBLE OR WEEDS, OR CUT FOR STOCKFOOD, FOR 4 WEEKS AFTER APPLICATION.

Sugarcane: DO NOT GRAZE CROP OR SPRAYED VEGETATION, OR CUT FOR STOCKFOOD, FOR 6 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

TITAN Imazapic 240 Herbicide is a member of the imidazolinone group of herbicides. The product has the inhibitors-of-acetolactate-synthase (ALS) mode of action. For weed resistance management, the product is a Group 2 herbicide. Some naturally-occurring weed biotypes resistant to this product and other Group 2 herbicides, may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population, if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

ALL APPLICATIONS

Reduce the incidence of "fines" by using the correct nozzles and operating pressures.

Replace worn or old nozzles and calibrate equipment on a regular basis.

Fallow application: TITAN Imazapic 240 Herbicide can be applied in fallow prior to the planting of various crops as outlined in the Follow Crops table. TITAN Imazapic 240 Herbicide is a pre-emergence herbicide, which is recommended to provide short term (6 to 12 weeks) pre-emergence control of summer germinating grasses and broadleaved weeds in cultivated or uncultivated fallow.

A green weed cover of greater than 50% will significantly reduce the length of pre-emergence weed control.

Sugarcane application

Pre-emergence: Where possible, apply to soil two days prior to incorporation. Incorporate with adequate irrigation or rainfall to a minimum depth of 5cm. Incorporation by flood irrigation may result in loss of weed control. TITAN Imazapic 240 Herbicide should be applied to plant or ratoon cane before the cane has emerged. When applying TITAN Imazapic 240 Herbicide in moist conditions, allow weeds to germinate prior to application and apply mixed with an appropriate knockdown herbicide. Whenever emerged weeds are present, TITAN Imazapic 240 Herbicide should be mixed with an appropriate knockdown herbicide.

Post-emergence: TITAN Imazapic 240 Herbicide MUST always be mixed with registered rates of paraquat and applied as a carefully directed inter-row spray to weeds and at the base of the sugarcane stalks, minimising spray contact with sugarcane foliage. DO NOT add crop oil to this mixture. Other surfactant or adjuvant is not required.

Peanut application

Pre-emergence: Apply to soil and incorporate with adequate irrigation or rainfall to a minimum depth of 5cm.

Post-emergence: TITAN Imazapic 240 Herbicide must always be mixed with an adjuvant as directed in the DIRECTIONS FOR USE section of the label.

SOIL TYPE

Pre-emergence weed control with TITAN Imazapic 240 Herbicide can be limited when applied onto soils with pH <5.0 and/or which contain high concentrations of iron (Fe) and/or aluminium (Al). These situations are common in soil types such as Krasnozems and red brown Ferrosols.

MIXING

TITAN Imazapic 240 Herbicide is an aqueous solution. Part fill the spray tank with water, then with agitator running, add the required amount of product, then fill the tank with water.

When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product. For post-emergence applications in peanuts, add adjuvant after the addition of TITAN Imazapic 240 Herbicide to the tank.

No adjuvants or surfactants are required when TITAN Imazapic 240 Herbicide is used alone for pre-emergence weed control or with paraquat.

DO NOT add crop oil when applied post-emergence with paraquat.

APPLICATION

DO NOT apply by aircraft.

Fallow application: Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and do not overspray headlands.

Sugarcane: Ensure thorough coverage of soil and weeds with well-calibrated spray equipment.

Apply in a minimum of 200L water/ha using a nozzle configuration to generate a medium to coarse spray pattern for pre-emergence applications or medium spray pattern for post-emergence applications. Replace worn or old nozzles and calibrate equipment on a regular basis.

GROUP 2 HERBICIDE

Peanuts: Apply with boom equipment in not less than 50L/ha water using flat fan nozzles.

Avoid overlap and do not overspray headlands. For post-emergence applications, TITAN Imazapic 240 Herbicide should be applied a minimum of 2 hours before rainfall.

EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of TITAN Imazapic 240 Herbicide and before use with other products.

COMPATIBILITY

TITAN Imazapic 240 Herbicide is compatible with Farmalinx PANDA 435, atrazine, glyphosate, paraquat, diquat, Ally*, Garlon*, Touchdown* and 2,4-D. It is NOT compatible with dicamba.

FOLLOW CROPS

Under conditions which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

Following use in summer fallow:-

MONTHS AFTER APPLICATION				
3	4	10	24	36
Faba Beans, Peanuts, Mung beans, Maize varieties with CLEARFIELD Technology ONLY: -Pacific hycorn 621T -Pacific hycorn 531T -Pioneer 33951R; Wheat varieties with CLEARFIELD Technology only	Wheat ¹ , (except for Wheat varieties with CLEARFIELD Technology ONLY), Barley ¹ , Lucerne ¹ , Chickpea	Sorghum ² , Maize ² (except for maize varieties with CLEARFIELD Technology ONLY), Dryland Cotton ³	Irrigated Cotton, Sunflower	any other crops

¹ The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), Barley or Lucerne during the next winter season.

DO NOT apply TITAN Imazapic 240 Herbicide in fallow later than the end of December.

DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200mm.

² The following additional requirements apply if it is intended to plant Sorghum or Maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of Sorghum or Maize is expected to be below 500mm.

³ The following additional requirements apply if it is intended to plant Dryland Cotton during the next summer season.

DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of dryland Cotton is expected to be below 550mm.

Following use in Sugarcane and Peanuts:-

MONTHS AFTER APPLICATION						
0	4	5	10	18	24	36
Mung beans, Peanuts, Soybeans, Maize varieties with CLEARFIELD Technology ONLY: -Pacific hycorn 621T -Pacific hycorn 531T -Pioneer 33951R	Chickpeas, Lucerne ¹ , Barley ¹ , Wheat ¹ (except for Wheat varieties with CLEARFIELD Technology ONLY)	Lupins, Pasture Legumes	Maize ² (except varieties with CLEARFIELD Technology ONLY), Dryland Cotton ³	Sorghum ²	irrigated Cotton, Sunflower	any other crops

¹ The following additional requirements apply if it is intended to plant wheat (except for varieties with CLEARFIELD Technology), Barley or Lucerne during the next winter season.



DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of cereals is expected to be below 200mm.

² The following additional requirements apply if it is intended to plant Sorghum or Maize (except for varieties with CLEARFIELD Technology) during the next summer season.

DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of Sorghum or Maize is expected to be below 500mm.

³ The following additional requirements apply if it is intended to plant dryland Cotton during the next summer season.

DO NOT apply TITAN Imazapic 240 Herbicide in areas where rainfall from spraying to sowing of dryland Cotton is expected to be below 550mm.

RE-ENTRY PERIOD

When tank-mixing TITAN Imazapic 240 Herbicide with Farmalinx PANDA, DO NOT allow entry into treated areas for 12 hours. When prior entry is necessary, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. DO NOT spray within 50m of wetlands or waterways.

CROP SAFETY

This product may, in some circumstances, lead to transient crop yellowing and temporary slowing of growth in Sugarcane but plants soon recover and yield is unaffected. This effect may be more pronounced when the product is used under conditions where soil moisture is high within 2 days following application and on light/sandy soil. Contact TITAN AG Pty Ltd for advice on variety susceptibility before using this product.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate streams, rivers, waterways or drains with this product or used containers.

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN. Store in the closed, original container in a dry, well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, puncture or shred and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray, wear elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves. When tank-mixing with Farmalinx PANDA or Paraquat, consult also the safety directions for those products.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Imazapic 240 Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Causes eye irritation. Precautionary Statements: Wash all exposed external body areas thoroughly after handling. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

* Other trademarks