

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

LV ESTER 680

HERBICIDE

ACTIVE CONSTITUENT: 680g/L 2,4-D-2-ETHYLHEXYL ESTER

GROUP **4** HERBICIDE

A specially formulated low volatile herbicide for selective control of various weeds in crops, pastures and non-agricultural areas as per the directions for use table.

THIS IS A PHENOXY HERBICIDE THAT CAN CAUSE SEVERE DAMAGE TO NATIVE VEGETATION AND SUSCEPTIBLE CROPS SUCH AS COTTON, GRAPES, TOMATOES, OILSEED CROPS AND ORNAMENTALS.

APVMA Approval No.: 63412/131557

Pack Size: 20L-1000L



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | Fax (02) 9999 0483
titanag.com.au

IN A TRANSPORT EMERGENCY

DIAL 000

POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL

DIRECTIONS FOR USE

General Restraints: DO NOT exceed maximum application rate of 6.6L/ha (4500g ae/ha).
DO NOT exceed the maximum daily application rate by backpack spraying of 5.9L/day.
DO NOT apply if heavy rains or storms are forecast within 3 days.
DO NOT irrigate to the point of run-off for at least 3 days after application.
DO NOT apply if crop or weeds are stressed due to dry or excessively moist conditions.

Additional USAGE restrictions apply in some crops, states and seasons, see restriction tables 1, 2, 3, 4 and 5.

Timing and Usage Restriction Tables

Table 1: Timing restrictions for spraying peanuts			
Situation	Rate (L/ha)	Region	Timing Restriction
			DO NOT APPLY DURING THE MONTHS
Broadcast spraying, prior to sowing (peanuts)	Up to 1.3L/ha	Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	October
		Mackay/Whitsunday	September to December
		Mary/Burnett	October to November
		SE Queensland	August to May
	Up to 1.6L/ha	Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	October
		Mackay/Whitsunday	August to December
		Mary/Burnett	September to November
		SE Queensland	Use not supported
Band spraying, post-sowing pre-emergence (peanuts)	Up to 1.6L/ha	Queensland dryland	No timing restrictions
		Cape York	No timing restrictions
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	No timing restrictions
		Burdekin	No timing restrictions
		Mackay/Whitsunday	No timing restrictions
		Mary/Burnett	No timing restrictions
		SE Queensland	October to January
Broadcast spray, post-sowing pre-emergence (peanuts)	Up to 3.3L/ha	Queensland dryland	June to August
		Cape York	October and November
		Northern Gulf	October and November
		Northern Territory	October and November
		Wet Tropics	October to December
		Burdekin	September and October
		Mackay/Whitsunday	August to December
		Mary/Burnett	April to January
		SE Queensland	Not supported

Table 2: Application and timing restrictions for application to pastures					
DO NOT apply above maximum rate (L/ha) below OR label rate, whichever is LOWEST.					
	State	Summer	Autumn	Winter	Spring
Pastures (prior to sowing, conservation tillage)	Queensland & NT	4.7	4.7	4.7	4.7
	New South Wales & ACT	4.7	4.7	4.7	4.7
	Victoria	0.5	1.5	4.7	1.5
	Tasmania	0.5	1.1	3.3	1.5
	South Australia	1.1	1.5	4.7	3.3
	Western Australia	1.5	3.3	4.7	3.3
	State	Summer	Autumn	Winter	Spring
Pastures (established)	Queensland & NT	6.6	6.6	6.6	6.6
	New South Wales & ACT	6.6	6.6	6.6	6.6
	Victoria	0.9	1.8	6.6	3.3
	Tasmania	0.6	1.5	4.7	2.9
	South Australia	1.3	2.9	6.6	4.7
	Western Australia	3.3	4.7	6.6	4.7

Table 3: Timing restrictions for spraying SUGARCANE		
Rate (L/ha)	Region	Timing Restriction
		DO NOT APPLY DURING THE MONTHS
Up to 1.2L/ha	All	No timing restriction
Up to 2.4L/ha	Wet Tropics	No timing restriction
	Burdekin	October
	Mackay/Whitsunday	September to December
	Mary/Burnett	August to December and April to May
	Northern NSW	No timing restriction

Table 4: Application restrictions for TURF		
DO NOT apply above maximum rate (L/ha) below OR label rate, whichever is LOWEST.		
	State	Rate (L/ha)
TURF	Queensland & NT	2.9
	New South Wales & ACT	2.9
	Victoria	2.3
	Tasmania	2.3
	South Australia	2.3
	Western Australia	3.7
If applying to golf courses in Tasmania, DO NOT apply to fairways adjacent to natural water bodies.		

Table 5: Risk mitigation measures for Dryland cropping, pre-emergent uses	
Situation	Risk mitigation measures
Dryland cropping, preparatory spray	Only apply in no-till farming systems (Tasmania, South Australia)
Winter cereals, pre-emergence uses	Only apply in no-till farming systems (Tasmania, South Australia, Western Australia)
Summer cereals, pre-emergent uses	Only apply in no-till farming systems (Tasmania, South Australia)

SPRAY DRIFT RESTRAINTS

DO NOT apply by a vertical sprayer.

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift

DO NOT allow **bystanders** to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an **unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production**, or cause contamination of plant or livestock commodities, outside the application site from **spray drift**. The buffer zones in the relevant buffer zone tables below provide guidance but may not be sufficient in all situations. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the **wind speed** is between 3 and 20 kilometres per hour at the **application site** during the time of application.

DO NOT apply if there are **hazardous surface temperature inversion** conditions present at the application site during the time of application. **Surface temperature inversion conditions** exist most evenings one to two hours before sunset and persist until one or two hours after sunrise.

BOOM SPRAYERS

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets not smaller than a VERY COARSE **spray droplet size category**.
- Minimum distances between the **application site** and downwind **sensitive areas** (see 'Mandatory downwind buffer zones' section of the following table titled 'Buffer zones for boom sprayers') are observed.

Buffer Zones for Boom Sprayers						
Application rate (ha)	Boom Height above target canopy	Mandatory buffer zones (distances given in metres)				
		Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to 800mL (560g ae/ha)	0.5m or lower	0 metres	10	0 metres	25	0 metres
	1.0m or lower		40		55	
Up to 1.7L (1150g ae/ha)	0.5m or lower		30		35	
	1.0m or lower		60		100	
Up to 2.4L (1620g ae/ha)	0.5m or lower		30		45	
	1.0m or lower		80		140	
Up to 4.7L (3180g ae/ha)	0.5m or lower		50		100	
	1.0m or lower		160		375	
Up to 6.6L (4500g ae/ha)	0.5m or lower		75		150	
	1.0m or lower		Not supported		Not supported	

AIRCRAFT

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a Very Coarse **spray droplet size category**.
- For maximum release height above the target canopy of 3 metres or 25% of wingspan or 25% of rotor diameter, whichever is the greatest, minimum distances between the application site and downwind sensitive areas (see 'Mandatory buffer zones' section of the following table titled 'Buffer zones for aircraft') are observed.

BUFFER ZONES FOR AIRCRAFT						
Application rate (ha)	Aircraft Type	Mandatory buffer zones (distances given in metres)				
		Bystander Areas	Natural Aquatic Areas	Pollinator Areas	Vegetation Areas	Livestock Areas
Up to 820mL (560g ae/ha)	Fixed Wing	0 metres	120	0 metres	170	0 metres
	Helicopter		85		120	
Up to 1.7L (1150g ae/ha)	Fixed Wing		190		300	
	Helicopter		130		190	
Up to 2.4L (1620g ae/ha)	Fixed Wing		240		400	
	Helicopter		160		240	
Up to 4.7L (3180g ae/ha)	Fixed Wing		Not supported		Not supported	
	Helicopter		275		400	
Up to 6.6L (4500g ae/ha)	Fixed Wing		Not supported		Not supported	
	Helicopter		350		625	

FIELD CROPS – REFER TO SECTION “SPRAY APPLICATIONS AND DRIFT RISK ASSESSMENT” BEFORE APPLICATION				
SITUATION AND CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Wheat, Barley	Refer to Weed Table	VIC only	210-800mL	CROP STAGES: ALL CEREALS Variations between varieties do occur. Check sensitivity and growth stages of varieties before applying. Damage may result if applied too early. VIC only: Apply at tillered to boot stages. NSW, ACT only: Apply after when the first node can be felt at the base of a tiller and before swelling of the head can be felt in a tiller. QLD only: Apply from mid-tillering (5 to 6 fully emerged main stem leaves plus one or more tillers) to before boot stage (visible swelling of the head at the top of the main stem). SA, TAS only: Apply from completion of tillering to early jointing stage. WA only: Apply from the 5 leaf stage up to jointing stage (Zadoks 15-33). Apply only at 6 leaf stage for Cranbrook and Jacup wheats (Zadok 16) to avoid possible damage.
		SA only	230-800mL	
		QLD, NSW, ACT only	410-800mL	
		TAS only	620-800mL	
		WA only	800mL	
Triticale	Refer to Weed Table	QLD, NSW, ACT only	410-800mL	
		SA only	240-820mL	
		VIC only	210-800mL	
Cereal Rye	Refer to Weed Table	QLD, NSW, ACT only	410-800mL	
		VIC only	210-800mL	
Sugarcane	Refer to Weed Table	QLD only	1.15-2.4L	Post-emergence.
Stubble/Fallow Spray Prior to Direct Drilling or Sowing. Winter Cereals, Grain Legumes (Peanuts QLD only), Canola	Refer to Weed Table	ALL STATES	210-800mL	Observe the plant back periods given in the table in this leaflet. Must be tank mixed with a knockdown herbicide such as TITAN Glyphosate 450 Herbicide, TITAN Paraquat 250 or TITAN EOS Herbicide. Select appropriate rate from the weed table. For Skeleton Weed, spraying should only be done 6-8 weeks before anticipated sowing date and subsequent cultivation limited to a minimum.
Harvest Aid or Salvage Spray	Broadleaf Weeds	ALL STATES	1.7L	Apply after dough stage of crop. Interval between application and effectiveness is 10-20 days. For desiccation of green matter, estimate harvest date and apply spray approximately 14 days earlier. Rain between spraying and actual harvest can negate results. Note: Where thistles are tall and branching above the crop, spraying can turn the branches down into the crop, presenting more stalks to cause header comb blockages. Spraying may increase seed contamination of harvest by accelerating maturity. DO NOT use with undersown legumes that have not set seed.
Winter Cereals	Refer to Weed Table			
Potatoes Pre-harvest Preparation	Broadleaf Weeds such as Clover, Variegated Thistle and Cruciferous weeds	VIC, TAS only	1.15-2.4L	Apply approximately 4 to 5 weeks before harvest after the potato haulms have dried off. Use the highest rate where weeds are more than 30cm in height. For boom spraying apply at least 100 litres of spray mixture per hectare. If grasses such as Rye Grass and Winter Grass are also present add Amitrole* T Herbicide.
PASTURES, NON-AGRICULTURAL AND INDUSTRIAL				
SITUATION AND CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Improved Pasture containing Clovers	Refer to Weed Table	QLD, NSW, ACT, TAS, SA only	410-800mL	Clover must be well covered by the grass or extensive damage may result.
Pastures – non legumes, Rights of Way & Industrial		QLD, NSW, ACT, TAS, SA only	800mL-4.7L	Control of most perennial weeds, but due to the rooting habits of most species control may take a number of years. Damage may result to legumes in pasture.
		VIC only	800mL-6.5L	Boom spray.
Pastures – Direct Drilling or Surface Sowing	Charlock, Clover, Medics, Mustards, Paterson’s Curse, Saffron, Slender, Variegated and Spear Thistles, Turnip Weed, Wild Radish, Wild Turnip As Above plus: Capeweed, Wireweed, Storksbill/Erodium, Flatweed, Horehound (seedlings), Skeleton Weed, Nodding or Star Thistles St John’s Wort All of above plus grasses	QLD, NSW, ACT, VIC, TAS, SA, WA only	800mL-1.5L (Aerial Application)	Apply to young, actively growing weeds. Sowing: DO NOT sow pasture seed for at least 21 days after application. If soil moisture is dry, delay sowing for at least 30 days.
			800mL-1.5L (Ground Application)	
			3.3-4.7L (Aerial or Ground)	
			As above plus Agricrop Propon* or TITAN Glyphosate 450 Herbicide	

SPOT SPRAYING				
SITUATION AND CROP	WEEDS CONTROLLED	STATE	RATE/ha	CRITICAL COMMENTS
Spot Spraying (All Situations)	Refer to Weed Table	ALL STATES	1/100th of rate on Weed Table per 10L water per 100m ²	Apply through Knapsack. Thorough wetting of weed is essential.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

IN TASMANIA, THIS PRODUCT MAY ONLY BE USED FROM 15TH APRIL TO 15TH SEPTEMBER UNLESS OTHERWISE PERMITTED BY THE REGISTRAR OF PESTICIDES.

WITHHOLDING PERIOD (WHP):

PASTURES, CEREAL CROPS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.

CROP HARVEST WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

WEED TABLE

Note: Where weeds are to be sprayed in a CROP or PASTURE, use only the rates given for the crop in the table below. In most cases this will give control, however some hard to kill weeds or those in advanced stages of growth may only be suppressed eg *Rumex* spp. (Docks) and *Polygonum* spp. (Wireweed, Climbing Buckwheat) are killed to ground level only.

WEEDS CONTROLLED	APPLICATION RATE PER HECTARE								CRITICAL COMMENTS
	Crop						Pasture - Non Legume		
	VIC	SA	TAS	NSW, ACT	QLD	WA	VIC	NSW, ACT, TAS, SA, QLD, WA only	
<i>Amaranthus</i> spp.	-	-	-	800mL	-	-	-	-	
Angled Onion	-	-	-	-	-	-	3.3L	800mL-1.7L	Spray when buds forming or early flowering.
Apple of Sodom	-	-	-	-	-	-	-	2.9-3.3L	
Bathurst Burr	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Black Knapweed	-	-	-	-	-	-	3.3L	-	Spray before flowering. DO NOT cultivate these infestations.
Buffalo Burr	-	-	-	-	-	-	-	800mL-1.15L (Not QLD & WA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
California Burr	-	-	-	800mL	-	-	1.7-3.3L	1.15-1.7L (Not SA)	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Caltrop	-	-	-	620-800mL	-	-	1.7-3.3L	-	Spray from seedling to pre-flowering. Use higher rate as plant matures.
Cape Tulip	-	-	-	-	-	1.15L	3.3L	1.7-3.3L	Spray before flowering.
Capeweed	800mL	800mL	800mL	530-800mL	-	-	-	2.5-3.3L	Spray up to rosette stage.
Charlock	410mL	410mL	800mL	410mL	-	-	-	800mL	Spray up to rosette stage.
Clover	-	-	-	620-800mL	-	-	-	800mL	
Colocynth	-	-	-	-	-	-	3.3L	-	Spray at seedling stage only.
Deadnettle	-	-	-	800mL	-	-	-	-	
Devil's Claw	-	-	-	800mL	-	-	1.3L	1.15-1.7L (Not SA)	Spray prior to pods forming.
Dock	800mL	800mL	-	-	800mL	800mL	-	1.7-2.5L	Spray at rosette stage to kill top growth only.
Fat Hen	-	-	-	410-800mL	-	-	-	-	
Flatweed	-	-	-	800mL	-	-	-	2.5-3.3L	
Fumitory (red)	-	800mL	-	800mL	-	-	-	2.5-3.3L	Spray up to rosette stage.
Fumitory (white)	800mL	410mL	-	800mL	-	-	-	2.5-3.3L	Spray up to rosette stage.
Galvanised Burr	-	-	-	-	-	-	4.7L	4.7L (Not QLD & WA)	Spray from seedling to pre-flowering.
Goosefoots	-	-	-	800mL	-	-	-	-	
Hard Head or Russian Knapweed	-	-	-	-	-	-	3.3-5.2L	-	Spray before flowering.
Hogweed, Wireweed	800mL	800mL	-	800mL	800mL	-	-	1.15-1.7L (Not SA)	Spray up to rosette stage.
Hoary Cress, Whiteweed	-	800mL	800mL	800mL	-	-	1.7-3.3L	1.7-2.1L	Spray from late rosette to pre-flowering.
Horehound (seedlings)	-	800mL	-	-	-	840mL	-	1.7-3.3L	Late autumn to early spring.
Ironweed, Corn Gromwell	-	-	-	800mL	-	-	-	1.15-1.7L	
Khaki Weed	-	-	-	-	-	-	-	800mL-1.15L (Not SA)	Spray young seedlings only.

WEEDS CONTROLLED	APPLICATION RATE PER HECTARE								CRITICAL COMMENTS
	Crop						Pasture - Non Legume		
	VIC	SA	TAS	NSW, ACT	QLD	WA	VIC	NSW, ACT, TAS, SA, QLD, WA only	
Lincoln Weed	-	800mL	-	-	-	-	-	-	Autumn spray before sowing improves control.
London Rocket	-	-	-	-	-	570mL	-	1.6-2.5L (WA only)	
Lupins	800mL	-	-	410-800mL	-	-	-	-	Spray up to rosette stage.
Melilotus/Hexham Scent	800mL	800mL	-	-	800mL	-	-	1.15-1.7L	Spray up to rosette stage.
Melons - Camel, Paddy	-	-	-	410-800mL	-	-	-	-	
Mustards	330mL	230-800mL	800mL	410-900mL	620mL	620mL	3.3L	1.7-2.5L	Spray up to rosette stage.
Mexican Poppy	-	2.3-3.5L	-	800mL	-	840mL	-	800mL-1.15L (1.1-1.5L WA only)	Spray rosette stage and before flowering.
Mintweed	-	-	-	800mL	620mL	-	-	800mL-1.15L	Spray active seedlings only.
Muskweed	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.
New Zealand Spinach	-	-	-	800mL	-	-	-	-	
Noogoora Burr	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	Spray seedling to pre-flowering.
Nut Grass	-	-	-	-	-	-	3.3-5.2L	-	Spray within 4 weeks of foliage emergence, repeat spray necessary.
Paterson's Curse	-	-	-	800mL	-	840mL	1.7-3.3L	800mL-1.7L (1.15-1.5L WA only)	Spray seedling to rosette stage.
Poppy Wild	410mL	-	-	-	-	-	-	2.1-2.9L	Spray up to rosette stage.
Ragwort	-	-	-	-	-	-	3.3L	3.3L	Spray at rosette to cabbage stage.
Rapeseed	800mL	-	-	410-800mL	-	-	-	-	Spray up to rosette stage.
Rapistrum	-	-	-	-	-	570mL	-	840mL (WA only)	
Rough Poppy	-	410mL	-	410-800mL	-	-	-	800mL	Spray young seedlings only.
St. John's Wort	-	-	-	-	-	-	3.3-5.2L	3.3-4.7L	Spray before flowering. Spray before plants are 40cm high.
Safflower	-	-	-	410-800mL	-	-	-	-	
Sand Mustard/ Sand Rocket	-	-	-	-	-	-	3.3L	-	Spray before flowering.
Shepherd's Purse	-	-	-	800mL	-	-	-	-	
Silver Leaf Nightshade	-	-	-	-	-	-	3.3L	-	Spray at flowering. Fallow land: controls top growth only.
Skeleton Weed	-	800mL	-	800mL	-	-	3.3L	1.15-1.7L	Spray rosettes before aerial growth commences.
Stingless Nettle (Deadnettle)	-	800mL	800mL	-	-	-	-	2.1-2.5L	
Stinging Nettle	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Stinkwort	-	-	-	800mL	-	-	1.7-3.3L	1.7-3.3L	Spray younger plants. Use higher rate as plants mature.
Storksbill/Erodium	-	-	-	800mL	-	-	-	-	
Sunflower Seedlings	800mL	-	-	410-800mL	620mL	-	-	-	Spray multiple leaves.
Thistles:									
- Golden	-	-	-	-	-	-	3.3L	3.3L	Spray at rosette stage.
- Nodding	-	-	-	-	-	-	3.3L	1.15-1.7L	Spray rosette stage to pre-flowering.
- Saffron	620mL	800mL	-	410-800mL	800mL	800mL	800mL-1.7L	800mL-3.3L (WA only)	Spray up to rosette stage.
- Sheep	-	-	-	-	-	840mL	-	840mL-3.3L (WA only)	
- Slender, Shore	-	-	800mL	800mL	-	-	1.7-3.3L	800mL-3.3L	Spray at rosette stage.
- Soldier	-	-	-	-	-	-	3.3L	-	Spray at rosette stage.
- Spear	800mL	-	800mL	-	-	-	800mL-2.5L	1.15-2.1L	Spray at seedling to rosette stage. Use higher rate as plants mature (pastures).
- Stemless	-	-	-	-	-	-	3.3L	2.5-3.3L	Spray rosette stage to flowering.
- St Barnaby's	-	-	-	-	-	-	-	1.15-1.7L	
- Star	-	-	-	800mL	-	-	1.7-3.3L	1.15-1.7L	Spray seedling to rosette stage. Use higher rate as plants mature.
- Variegated	-	-	800mL	410-800mL	620mL	-	800mL-2.5L	800mL-3.3L	Spray at rosette stage. Can cause stock poisoning.
Thornapple	-	3.5L	-	410-800mL	-	-	3.3L	800mL-1.7L	Spray at seedling stage.
Tree Hooweed	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.

WEEDS CONTROLLED	APPLICATION RATE PER HECTARE								CRITICAL COMMENTS
	Crop						Pasture - Non Legume		
	VIC	SA	TAS	NSW, ACT	QLD	WA	VIC	NSW, ACT, TAS, SA, QLD, WA only	
Turnip Weed	-	410mL	-	410-800mL	410mL	620mL	-	800mL	Spray seedlings only.
Vetches/Tares	800mL	620mL	800mL	-	-	-	-	-	
Wards Weed	-	410mL	-	-	-	-	-	-	Spray at seedling stage.
Wild Cabbage	800mL	-	-	-	-	-	-	-	Spray up to rosette stage.
Wild Garlic	-	-	-	-	-	-	6.6L	-	Suppresses aerial growth only.
Wild Mignonette	-	-	-	-	-	840mL	3.3L	-	Spray at rosette stage.
Wild Mustard	-	-	-	-	-	570mL	-	1.6-2.5L (WA only)	
Wild Radish	800mL	800mL	800mL	410-800mL	800mL	570mL	-	800mL (840mL WA only)	Spray up to rosette stage.
Wild Sage	-	-	-	-	-	-	-	2.5-3.3L	
Wild Teasel	-	-	-	-	-	-	1.7-3.3L	-	Spray at rosette stage. Use higher rate as plants mature.
Wild Turnip	210mL	230mL	800mL	410-800mL	-	400mL	-	800mL (840mL WA only)	Spray up to rosette stage.

Plant back days for TITAN LV ESTER 680 HERBICIDE			
CROP	RATES		
	Up to 510mL/ha	510mL-1L/ha	1-1.6L/ha
Balansa Clover	7	7	10
Barley ¹	1	1	3
Chickpeas ²	7	14	21
Cotton	10	14	21
Faba Beans	7	7	10
Field Peas	7	14	14
Lentils	7	7	10
Linseed	7	7	14
Lucerne	7	7	10
Lupins ⁴	7	14	21
Medics	7	7	10
Narbon Beans	7	7	10
Navy Beans	10	10	14
Oats	3	3	7
Perennial Ryegrass	7	7	10
Persian Clover	7	7	10
Phalaris	7	7	10
Canola/Rapeseed ²	14	21	28
Rice	7	7	14
Safflower ²	7	14	21
Sorghum ³	3	7	10
Soybean	14	14	21
Sub-Clover	7	7	10
Sunflower ³	7	10	14
Triticale ¹	1	3	7
Vetch	7	7	10
Wheat ¹	1	3	7
White Clover	7	7	10

IMPORTANT: WHEN APPLIED TO DRY SOILS AT LEAST 15mm (1/2 inch) OF RAIN MUST FALL PRIOR TO THE COMMENCEMENT OF THE PLANT BACK PERIOD.

Notes:

1. In Queensland, no rainfall is required to fall prior to commencement of Plant Back Period for Wheat, Barley and Triticale.
2. In Queensland, planting of Canola/Rapeseed, Chickpeas and Safflower must be delayed for at least 14 days following rainfall at least 15mm.
3. In Central Queensland, when using 730mL/ha or less of TITAN LV Ester 680 Herbicide, the Plant Back Period for Sorghum and Sunflower is 1 day irrespective of rainfall.
4. In WA the Plant Back Period for Lupins at all rates is 28 days.

GENERAL INSTRUCTIONS

Before opening, carefully read Directions For Use, Precautionary Statements, Safety Directions and First Aid Instructions.

APPLICATION INFORMATION

This product may be used in either high or low volume sprays. Just pour into water and stir.

Boom Spraying: Use 30-100 litres water per hectare.

Aerial Spraying: Use 40-90 litres water per hectare.

Note: Refer to the Department of Agriculture/Primary Industries in your state for the current restricted spraying areas.

EQUIPMENT MAINTENANCE AND USAGE

Keep the spray unit for herbicides only if possible. Otherwise wash out the unit with hot soapy water followed by several clear water rinses. DO NOT use wooden spray vats as they can not be cleaned. Hoses cannot be cleaned and new hoses should be fitted when the unit is to be used for any other purpose.

COMPATIBILITY

This product can be tank mixed with TITAN Dicamba 500 Herbicide, TITAN Chlorsulfuron 750 WG Herbicide, TITAN Paraquat 250 Herbicide, Agricrop Propon*, TITAN Atrazine 900 WG Herbicide, TITAN Glyphosate 450 Herbicide, TITAN Metsulfuron 600 WG Herbicide, TITAN Triasulfuron 750 WG Herbicide, TITAN EOS Herbicide, TITAN Triclopyr 600 Herbicide, TITAN Fluroxypyr 200 EC Herbicide.

TANK MIXING INSTRUCTIONS

- Fill the spray tank 1/4 full with water and agitate.
- Add wettable powders and water dispersible granules first.
- Agitate until these are uniformly dispersed, while adding water until the tank is 90% full.
- Add suspension concentrates (flowables) then soluble concentrates. Add emulsifiable concentrates last.
- Top up the tank with water and continue agitation until all the ingredients are properly mixed.
- Observe any mixing sequence instructions specifically stated on the tank mix products.

RESISTANT WEEDS WARNING

TITAN LV Ester 680 Herbicide is a member of the Phenoxys group of herbicides. TITAN LV Ester 680 Herbicide has the Disruptors of plant cell growth mode of action. For weed resistance management TITAN LV Ester 680 Herbicide is a Group 4 herbicide. Some naturally-occurring weed biotypes resistant to TITAN LV Ester 680 Herbicide and other Group 4 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by TITAN LV Ester 680 Herbicide or other Group 4 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN LV Ester 680 Herbicide to control resistant weeds.

RE-ENTRY PERIOD

DO NOT hand harvest Sugarcane for at least 1 day after application. If re-entering treated areas before the spray has dried, workers should wear overalls, elbow-length gloves and water-resistant footwear. Clothing must be laundered after each day's use.

PROTECTION OF CROPS, NATIVE AND NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

INTEGRATED PEST MANAGEMENT

Toxic to beneficial arthropods. Not compatible with integrated pest management (IPM) programs utilising beneficial arthropods. Minimise spray drift to reduce harmful effects on beneficial arthropods in non-crop areas.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty

packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

110L: Store in the original sealed Envirodrum in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT tamper with the Micro Matic valve or the security seal. DO NOT contaminate the drum with water or any water or any other foreign matter. After each use of the product, please ensure the Micro Matic coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained accordingly. When the contents of the drum have been used, please return the empty drum to the point of purchase. The drums remain the property of TITAN AG Pty Ltd.

Refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. When opening the container and preparing spray or using undiluted concentrate, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and half face piece respirator with organic vapour/gas cartridge or canister. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length chemical resistant gloves. If applying by hand wear half facepiece respirator with organic vapour/gas cartridge or canister. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, goggles, respirator (and if rubber wash with detergent and warm water) and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN LV Ester 680 Herbicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

* Other trademarks

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Harmful if swallowed. May cause an allergic skin reaction. Toxic to aquatic life with long lasting effects. Precautionary: Avoid breathing dust/fume/gas/mist/vapours/spray. Wear protective gloves/protective clothing/eye protection/face protection. Avoid release to the environment. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Specific treatment (see on this label). IF SWALLOWED: Call a POISON CENTER/doctor if you feel unwell. If skin irritation or rash occurs: Get medical advice/attention. Rinse mouth. IF ON SKIN: Wash with plenty of water. Collect spillage. Take off contaminated clothing and wash it before reuse. Dispose of contents/container in accordance with local/regional/national regulations.

