

# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

## BIFENTHRIN 250 EC INSECTICIDE

ACTIVE CONSTITUENT: 250g/L BIFENTHRIN  
SOLVENT: 666g/L HYDROCARBON LIQUID

GROUP **3A** INSECTICIDE

Controls insect pests and mites of apricots, bananas, barley, canola, citrus, clover, cotton, faba beans, field peas, grapes, lucerne, lucerne seed crops, lupins, navy beans, nectarines, peaches, pears, plums, subterranean clover, sugarcane, tomatoes and wheat as specified in the Directions for Use table.

APVMA Approval No.: 88918/122895

Pack Size: 1L; 5L; 10L; 20L; 200L



TITAN AG Pty Ltd | ABN 57 122 081 574  
15/16 Princes Street, Newport NSW 2106  
Tel (02) 9999 6655 | Fax (02) 9999 0483  
titanag.com.au

IN A TRANSPORT EMERGENCY

**DIAL 000**

POLICE OR FIRE BRIGADE



UN 3352  
PYRETHROID PESTICIDE,  
LIQUID, TOXIC (Bifenthrin  
(ISO)), ENVIRONMENTALLY  
HAZARDOUS  
PG: III Hazchem: 2X

## DIRECTIONS FOR USE

**Restraints:** DO NOT use as a foliar spray in banana plantations and orchards where mite predators or other beneficials are established and providing effective mite control and/or other pest control.  
DO NOT apply as a foliar treatment if rainfall is expected before spray deposits dry on leaf surfaces.  
DO NOT apply to bananas by aircraft.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Bananas	Banana Weevil Borer, Banana Rust Thrips	QLD, NSW, WA, NT only	<b>Seasonal Program</b> <u>Stool Treatment Method</u> 100-130mL/100L twice per year OR 265mL/100L once per year <u>Band Treatment Method</u> 100mL/100L twice per year <b>Monitoring Program</b> <u>Stool Treatment Method</u> 130mL/100L <u>Band Treatment Method</u> 100mL/100L	1 day	<b>Seasonal Program</b> <u>Twice per year Timing</u> Apply in October/November (spring/early summer) and March/April (late summer/autumn). Use the higher rate (concentration) when borer pressure or damage is high. <u>Once per year Timing</u> Apply in October/November OR March/April. <b>Monitoring Program</b> Monitor Weevil Borer populations carefully by trap counts and/or corm damage ratings, beginning in September when pest activity is on the increase and continue until April. Apply treatment when banana weevil borers reach or exceed acceptable threshold levels. Monitor borer control after application and re-treat as required. <b>Banana Weevil Borer:</b> Application should be made after rain or irrigation during periods of high adult borer activity. <b>Banana Rust Thrips:</b> Application against Banana Weevil Borer will give coincident rust thrips control, particularly when application is made when thrips activity is on the increase usually beginning September and into the summer months. <b>Application Method</b> <u>Stool Treatment Application</u> Remove trash from the base of stools and apply 500-750mL of spray solution to each stool, depending on stool size. Treat the bottom 30cm of each stool as well as the soil in a 30cm band around each stool, ensuring thorough treatment of both butt(s) and follower(s). Use the lower spray volume of 500mL on small stools less than 50cm across the entire base. <u>Band Treatment Application</u> Apply as a band application with a side delivery boom and offset nozzles on both sides of the row with the spray pattern positioned to spray 30cm of soil on either side of the row and 30cm in height. Aim to apply a total spray volume of 1L/stool area. For single sucker row configurations apply 28L of solution per 100 metres of row in a band 0.5m wide on each side of the row overlapping in the centre. For double sucker row configurations apply 56L of solution per 100 metres of row in a band 1m wide on each side of the double row with the spray pattern overlapping between the rows.
Bananas	Strawberry Spider Mite	QLD, WA only	16mL/100L	8 days	Monitor mite population on old leaves particularly during hot dry conditions. Apply TITAN Bifenthrin 250 EC Insecticide as a preventative rather than a curative treatment before damage occurs, and before mite numbers build up to damaging levels. Follow up applications may be required at 10-14 day intervals. Thorough coverage of the lower leaf surface is essential to ensure good control. Use a total spray volume of 300-500L/ha.
Cotton	Native Budworm, Cotton Bollworm, Two-spotted Mite, Green Mirid, Apple Dimpling Bug	QLD, NSW, WA only	240-320mL/ha	14 days (H) DO NOT graze or cut for stockfeed. DO NOT feed cotton trash to livestock.	Apply as indicated by field checks. Use the higher rate when pest pressure is high, conditions favour pest development and when increased residual protection is required. <b>Budworm and Bollworm:</b> Applications should be timed to coincide with egg hatch and when small larvae up to 5mm are present. DO NOT apply this product to cotton bollworm larvae larger than 5mm in length. <b>Two-spotted Mite:</b> Applications against Budworm and Bollworm will give good control of coincident Two-spotted Mite, particularly when applied on low mite populations (around 10% leaf infestation). If conditions continue to favour mite development a second application may be required 14-20 days later. <b>Green Mirid and Apple Dimpling Bug:</b> Apply at recommended threshold levels as indicated by field checks. Use the higher rate for increased pest pressure and longer residual protection.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Cotton – <i>continued</i>	False Wireworm, Sugarcane Wireworm	QLD, NSW, WA only	150mL/ha* or 1.5mL/100m of row	14 days (H)  DO NOT graze or cut for stockfeed. DO NOT feed cotton trash to livestock.	<b>Wireworms:</b> Apply as a spray into the furrow at planting. Use a spray nozzle which will deliver a coarse spray in a total volume of 60-100L/ha in a 10cm band over the seed before soil is brought in behind covering tynes in front of the press wheel.  * The rate is based on a 1m row spacing. If row spacing varies from 1m then apply at the use rate according to mL/100m of row.
Canola, Faba beans, Subterranean clover, Clover, Barley, Field peas, Lupins, Lucerne, Wheat	Redlegged Earth Mite, Brown Pasture Looper	ALL STATES	20 to 40mL/ha	4 weeks (G)	Apply as a broadcast ground rig application in a total water volume of 50-200L/ha or by air in a minimum total water volume of 20L/ha. Apply to bare soil after conventional cultivation and sowing or onto well grazed or sprayed pasture after direct drilling. Treat infested paddocks after sowing and before or soon after seedling emergence. Use the higher rate on heavier infestations and for longer residual protection. TITAN Bifenthrin 250 EC Insecticide is compatible with some herbicides. See compatibility statement for details. Use the 40mL rate when pest pressure is low. Monitor adjacent habitat and edges of the field for the presence of Vegetable Weevil prior to making a decision whether to spray.
	Blue Oat Mite, Pasture Webworm		40mL/ha		
	Bryobia Mites		80mL/ha		
Canola	Vegetable Weevil	ALL STATES	40-80mL/ha	4 weeks (G)	
Citrus	Leaf Eating Weevil	ALL STATES	  <b>Pre-emergence program</b> 5 or 10mL/tree  <b>Post-emergence monitoring program</b> 2.4mL/tree	–	Apply as a high volume band application in a 1.5 to 2 metres wide swath, to the ground, both sides of the row, under each tree. Aim to apply a total spray volume of 5 to 10L/tree (e.g. at 250 trees/ha = 1250 to 2500L/ha).  <b>Pre-emergence program:</b> Apply just prior to, or at the first sign of major beetle emergence in mid-October. Use the higher rate in blocks with a history of high beetle numbers or when longer residual control is required.  <b>Post-emergence monitoring program:</b> Apply at peak beetle emergence in October/November as indicated by field monitoring. (Refer to monitoring statement on label). Follow up treatment may be necessary based on a threshold of 25 beetles per 10 sites per orchard in consecutive counts 1-2 weeks apart.
Grapes	Fig Longicorn	NSW, ACT, WA only	400mL/100L	–	The application <b>MUST</b> be made at late dormancy after pruning and before bud burst. Apply a single high volume spray, with nozzles directing the spray solution to the trunk and cordons (arms) of grape vines to achieve thorough wetting of the bark. Total spray volume should be about 500mL/vine achieved by hand application.
Lucerne seed crops	Native Budworm	ALL STATES	160-240mL/ha	–	DO NOT treat lucerne seed crops for alfalfa sprout production. Apply as indicated by field checks after the commencement of flowering. Use the higher rate when pest pressure is high, conditions favour pest development and when increased residual protection is required.  <b>Native Budworm:</b> Applications should be timed to coincide with egg hatch and when small larvae up to 5mm are present.
Navy Beans	Native Budworm, Corn Earworm	ALL STATES	240-320mL/ha	14 days (H, G)	Apply as indicated by field checks from flowering onwards. Use the higher rate when pest pressure is high, conditions favour pest development and when increased residual protection is required.  <b>Budworm and Earworm:</b> Applications should be timed to coincide with egg hatch and when small larvae up to 5mm are present. DO NOT apply this product to corn earworm larvae larger than 5mm in length.
Peaches, Nectarines, Plums, Apricots	Carpophilus Beetle	ALL STATES	<b>Dilute spraying</b> 20mL/100L  <b>Concentrate spraying</b> Refer to the Mixing/ Application section	1 day	Monitor stone fruit orchards for Carpophilus Beetle as fruit approach maturity and become susceptible to attack. Apply TITAN Bifenthrin 250 EC Insecticide as a dilute spray before beetles reach damaging levels. Apply to the foliage and fruit of trees. Continue to monitor beetle numbers and if necessary reapply TITAN Bifenthrin 250 EC Insecticide up to 1 day before harvest or use another insecticide registered for this purpose. Apply no more than 2 applications per season. <b>There must be a minimum of 10 days between the re-treatment and the initial application.</b> Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. DO NOT use at rates greater than 40mL per 100L of water when using concentrate spraying. Cultural control methods (eg. destruction of fallen fruit by mulching) should be used to prevent excessive build up of Carpophilus Beetle.

CROP	PEST	STATE	RATE	WHP	CRITICAL COMMENTS
Pears	Longtailed Mealybug	VIC, WA only	10mL/100L plus Ampol DC Tron at 1L/100L	14 days	Examine wood for the presence of over wintering Longtailed Mealybugs but do not spray until large numbers of young nymphs emerge in spring. Apply this mixture to near the point of runoff to all above ground parts of the tree between green tip to commencement of flowering. DO NOT spray after flowering has commenced.
Sugarcane	Sugarcane Wireworm	QLD, NSW, WA only	150mL/ha* or 2.2mL/100m of row	–	Apply as a spray into the furrow at planting. Use a spray nozzle which will deliver a coarse spray in a total volume of 60-100L/ha in a band 20-30cm wide over the base of the furrow on top of the setts and before covering soil is brought in by tynes. * The rate is based on 1.5m row spacing. If row spacing varies from 1.5m then apply at the use rate according to mL/100m of row.
Tomatoes	Native Budworm, Corn Earworm, Two-spotted Mite, Tomato Russet Mite	ALL STATES	<b>High Volume</b> 16-24mL/100L or <b>Low Volume</b> 240mL/ha	1 day	DO NOT use low volume ground or air application on trellis tomatoes. <b>Crop Monitoring Program</b> <b>Budworm and Earworm:</b> Apply as indicated by field checks. Applications should be timed to coincide with egg hatch and when small larvae up to 5mm are present. DO NOT apply this product to corn earworm larvae larger than 5mm in length. <b>Mites:</b> Applications against Budworm and Earworm will give good control of coincident mites, particularly when applied on low mite populations. If conditions continue to favour mite development, a second application may be required 14-20 days later. <b>Schedule Spray Program</b> If fields are not checked during pest infestation periods, apply on a 7-10 day alternating program with a non-pyrethroid insecticide. Use the higher rate (high volume application) and shorter interval when pest infestation is more severe and when increased residual protection is required. DO NOT apply this product to Corn Earworm larvae larger than 5mm in length.
	Whitefly		12mL/100L water		Apply as indicated by pest incidence and repeat as necessary. Use a total spray volume of 2500L/ha.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS:**

**Tomatoes, Peaches, Nectarines, Plums, Apricots: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.**

**Bananas:**

**For Ground Applications: DO NOT HARVEST FOR 1 DAY AFTER APPLICATION.**

**For Foliar Applications: DO NOT HARVEST FOR 8 DAYS AFTER APPLICATION.**

**Cotton: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

**DO NOT GRAZE OR CUT FOR STOCKFEED.**

**DO NOT FEED COTTON TRASH TO LIVESTOCK**

**Pears: DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.**

**Navy Beans: DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.**

**Canola, Subterranean Clover, Clover, Field Peas, Faba Beans, Wheat, Barley, Lucerne, Lupins: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.**

**HARVEST WHP: NOT REQUIRED WHEN USED AS DIRECTED.**

**Citrus, Grapes, Sugarcane: NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

TITAN Bifenthrin 250 EC Insecticide can be used as a protective treatment when applied at regular intervals or as a knockdown treatment to control existing pests. Best results are obtained when TITAN Bifenthrin 250 EC Insecticide is applied before pest populations build up to damaging levels. This product is not suitable for use in Integrated Pest Management (IPM) programs where mite or other insect predators or parasites are established and providing effective mite and other insect control.

## APPLICATION

TITAN Bifenthrin 250 EC Insecticide may be applied by either ground rig or aircraft. Thorough coverage is essential to ensure adequate control. DO NOT apply as a fog or mist.

### Dilute Spraying:

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

### Concentrate Spraying:

- a) Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- b) Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- c) Determine an appropriate dilute spray volume (See Dilute Spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- d) The mixing rate for concentrate spraying can then be calculated in the following way:

Example only

1. Dilute spray volume as determined above: For example 1000L/ha.
2. Your chosen concentrate spray volume: For example 500L/ha.
3. The concentration factor in this example is:  $2 \times$  (ie.  $1000L \div 500L = 2$ ).
4. If the dilute label rate is 50mL/100L, then the concentrate rate becomes  $2 \times 50$ , that is 100mL/100L of concentrate spray.
5. The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
6. For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**Ground Application:** Applications should be made as a fine spray quality preferably using hollow cone nozzles. The application volume will depend on the type of crop to be treated. The following are suggested:

Low volume broadcast applications to – e.g. Cereals, Canola, Grain Legumes, Lucerne, Subterranean Clover: 50-200L/ha.

Low volume row crops applications to Cotton, Tomatoes, Navy Beans: 50-200L/ha.

High volume applications to row crops – e.g. trellised Tomatoes: 200-1500L/ha except as noted in critical comments. Use 200L/ha from transplanting increasing to 1500L/ha at maturity.

### High volume directed spray:

**Grapes:** Apply by hand application, using a high volume coarse spray of 500mL/vine. (e.g. at approx. 2500 vines/ha = 1250L/ha).

Foliar sprays to Bananas: 300 to 500L/ha.

High volume application to stone fruit: 1000 to 2000L/ha.

### Soil Applied Sprays:

#### High volume application

##### Bananas:

Stool treatment: Apply as a coarse spray at 500-750mL per stool.

Band treatment: Apply as a band application with a side delivery boom and offset nozzles – 1L of spray solution per stool.

**Citrus:** Apply as a high volume, directed spray to the ground under each tree. For optimum control apply to both sides of the tree. Total spray volume should be 5 to 10L/tree (e.g. at 250 trees/ha = 1250 to 2500L/ha).

### In furrow applications:

**Cotton & Sugarcane:** Use a coarse spray: 60 to 100L/ha as a band over the seed or sett before covering with soil – refer to critical comments for details.

## Aerial Application:

Use at least 20L/ha of total spray volume. Spray during the cooler parts of the day or night. To reduce possibility of drift, avoid spraying in calm conditions or when wind is light and variable. Preferably, spray in a crosswind. Use suitable application equipment and/or nozzles to deliver a fine spray quality. A spraydrift minimisation strategy should be employed at all times when aerially applying sprays to, or near, sensitive areas. The strategy envisaged is best exemplified by the cotton industry's Best Management Practice manual.

## MONITORING

**Post-emergence monitoring of Citrus Leaf Eating Weevil populations:** At first sign of major beetle emergence in mid October commence monitoring at 1 to 2 week intervals. Place polystyrene fruit box (330 x 480mm) under tree, shake branches vigorously, repeat on ten randomly selected trees throughout orchard. If 25 beetles or more are recorded in consecutive counts, treatment is required.

## MIXING

Add the required quantity of TITAN Bifenthrin 250 EC Insecticide to water in the spray tank and mix thoroughly. Maintain agitation during mixing and application.

## COMPATIBILITY

TITAN Bifenthrin 250 EC Insecticide is compatible with commonly used fungicides such as Dithane M45<sup>+</sup>, Antracol<sup>+</sup>, Barrack<sup>+</sup>, Bravo<sup>+</sup> and the herbicides – Spray.seed<sup>+</sup>, Broadstrike<sup>+</sup>, Spinnaker<sup>+</sup>, Simagranz<sup>+</sup>, Dual<sup>+</sup>, Sencor<sup>+</sup>, Glean<sup>+</sup>, Logran<sup>+</sup> and Stomp<sup>+</sup>.

## SURFACTANTS

TITAN Bifenthrin 250 EC Insecticide contains a surfactant. Additional surfactant may only be necessary on hard to wet plants and in high volume situations.

### \* NOTICE \*

*Helicoverpa* (= *Heliothis*) *armigera* resistance in Northern NSW and QLD. To help contain pyrethroid resistance in *H. armigera*, the Summer Crop Insecticide strategy as developed by the QLD Department of Agriculture and Fisheries and NSW Department of Primary Industries should be adhered to. Failure to observe the strategy may result in widespread resistance affecting the future viability of summer cropping.

## INSECTICIDE RESISTANCE WARNING

For insecticide resistance management TITAN Bifenthrin 250 EC Insecticide is a

GROUP	<b>3A</b>	INSECTICIDE
-------	-----------	-------------

Group 3A insecticide. Some naturally occurring insect biotypes resistant to TITAN Bifenthrin 250 EC Insecticide and other group 3A insecticide may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if TITAN Bifenthrin 250 EC Insecticide or other group 3A insecticides are used repeatedly. The effectiveness of TITAN Bifenthrin 250 EC Insecticide on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Bifenthrin 250 EC Insecticide to control resistant insects. TITAN Bifenthrin 250 EC Insecticide may be subject to specific resistance management strategies. For further information, contact your local supplier, TITAN AG representatives or local agricultural department agronomist.

## STONE FRUIT EXPORT ADVICE

Export of Treated Stone Fruit – Some export markets do not have suitable Maximum Residue Limits or import tolerances in place. Please contact TITAN AG Pty Ltd or the Australian Fresh Stone Fruit Growers Association prior to using this product on crops destined for export.

## RE-ENTRY TO TREATED FIELDS/CROPS

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

## PRECAUTION

DO NOT use human flaggers/workers unless they are protected by engineering controls such as enclosed cabs.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Dangerous to fish and aquatic organisms. DO NOT contaminate streams, rivers or waterways with the product or the used containers. Tail drains which flow from treated areas should be prevented from entering river systems.

## PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Spray in the early morning when bees are not actively foraging.



## STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area and away from children, animals, food, feedstuffs, seeds and fertilisers. DO NOT store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

**Refillable containers:** Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

## SAFETY DIRECTIONS

Attacks eyes. Poisonous if swallowed. Harmful if inhaled. Will irritate the skin. Avoid contact with eyes and skin. DO NOT inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, and elbow-length PVC gloves and goggles. If applying by hand, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat and elbow length PVC gloves. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If swallowed, do NOT induce vomiting.

## SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Bifenthrin 250 EC Insecticide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit [www.titanag.com.au](http://www.titanag.com.au)

**CONDITIONS OF SALE:** TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: Combustible liquid. Fatal if swallowed. Harmful if inhaled. May cause an allergic skin reaction. Suspected of causing cancer. Causes damage to the nervous system through prolonged or repeated exposure. May be fatal if swallowed and enters airways. Very toxic to aquatic life with long lasting effects. Precautionary Statements: Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Keep away from flames and hot surfaces. No smoking. Do not breathe dust/fume/gas/mist/vapours/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in a well-ventilated area. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF SWALLOWED: Immediately call a POISON CENTER/ doctor. Specific treatment (see on this label). Rinse mouth. Do NOT induce vomiting. IF ON SKIN: Wash with plenty of water. IF INHALED: Remove person to fresh air and keep comfortable for breathing. IF exposed or concerned: Get medical advice/attention. Call a POISON CENTER/doctor if you feel unwell. Get medical advice/attention if you feel unwell. Take off contaminated clothing and wash it before reuse. If skin irritation or rash occurs: Get medical advice/attention. In case of fire: Use to extinguish: CO<sub>2</sub>, powder or water spray. Collect spillage. Store in a well-ventilated place. Keep cool. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations.

