

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

TITAN

MANCOZEB 750 WG

FUNGICIDE

ACTIVE CONSTITUENT: 750g/kg MANCOZEB

GROUP **M3** FUNGICIDE

For the control of certain fungus diseases of fruit, field crops, tobacco, turf, vegetables and ornamentals
as per the Directions for Use Table.

APVMA Approval No.: 62383/120286

Pack Size: 20-25kg



TITAN AG Pty Ltd | ABN 57 122 081 574
15/16 Princes Street, Newport NSW 2106
Tel (02) 9999 6655 | Fax (02) 9999 0483
titanag.com.au

IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE

TRANSPORT AND HANDLING
NOT A DANGEROUS GOOD
ACCORDING TO THE AUSTRALIAN
DANGEROUS GOODS (ADG) CODE
FOR TRANSPORT BY ROAD AND RAIL

DIRECTIONS FOR USE

For use In ALL STATES where appropriate for crop and/or disease.

Restraint: DO NOT incorporate this product with an Integrated Mite control programme.

| TREE AND VINE CROPS | | | | | |
|---|---|---|---|---|--|
| RATE | | | | CRITICAL COMMENTS | |
| In the following table, all rates are given for dilute spraying. For concentrate spraying refer to the application section. | | | | For all uses in this table: Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods. | |
| CROPS | DISEASE/PEST | RATE | | WHP DAYS Harvest (H) Grazing (G) | |
| Citrus | Black Spot | 200g/100L plus spray oil (used according to its label instructions) | | Not required | On heavy soil: Spray at 8 and 15 weeks following a Bordeaux spray (5-5-100) at petal fall. On light soil: Add an adjuvant (spreader/sticker) according to label recommendations per 100L. Spray at 6 and 13 weeks following a Bordeaux spray (3.5-3.5-100) at petal fall. QLD, NT: Spray at 6 and 12 weeks after a copper spray at 1/2 to 3/4 petal fall. Apply at first sign of mite activity on fruit, usually between December and May. Choose a spray oil recommended for use on citrus. Apply when blossom malformation is observed, usually February. Choose a spray oil recommended for use on citrus. |
| | Citrus Rust Mite, Brown Citrus Mite | | | | |
| | Citrus Bud Mite | | | | |
| Custard Apples | Pseudocercospora Fruit Spot | 200g/100L | | 1 day (H) | DO NOT apply during flowering. Apply at 3-4 week intervals from first fruit set until harvest. Apply at 2 week intervals in wet, cloudy weather. |
| Grapevines | Black Spot, Downy Mildew | 200g/100L | | 30 days (H) | For Black Spot control apply every 10-14 days starting at bud burst. If Downy Mildew is expected, continue spray programme at intervals of 10-14 days until the threat of disease has passed. Use the shorter spray interval during weather conditions favourable to disease development. Apply at budburst and then repeat 7 to 10 days later. |
| | Phomopsis Cane and Leaf Spot | 150 to 200g/100L | | | |
| Mangoes | Anthrachnose (<i>Collectotrichum gloeosporioides</i>) | 200g/100L | | 1 day (H) | Spray at weekly intervals during flowering then monthly until nearer harvest. |
| Passionfruit | Septoria Leaf Spot, Brown Spot (<i>Alternaria passiflorae</i>) | 150g/100L | | 1 day (H) | Apply at 10-14 day intervals from October to May and every 21-28 days for remainder of year. Spray at weekly intervals during flowering and then fortnightly until near harvest. |
| | Anthrachnose (NSW only) | 200g/100L + 600mL/100L miscible summer oil | | | |
| Pawpaw (Papaya) | Black Spot | 200g/100L | | 1 day (H) | Apply with a wetting agent at 10 to 14 day intervals during periods conducive to disease. |
| Pome fruit: Apples and Pears | Apple Scab (Black Spot), Bitter Rot, Target Spot, Fly Speck, Sooty Blotch | 150 to 200g/100L | | 14 days (H) | Apply at 7-14 day Intervals following a cover copper spray at green tip. Use higher rate or shorter interval between sprays where conditions favour disease. This product can be mixed with Systhane 400WP Fungicide for the control of Apple Scab (Black Spot). Read and follow all the recommendations on the label of the tank mix partner. This product may be harmful to <i>Typhlodromus pyri</i> and its use in orchards where Integrated control is practiced should be avoided. |
| Stone Fruit: Almonds, Apricots, Cherries, Nectarines, Peaches, Plums* | Brown Rot, Rust, Shot Hole, Freckle | 150 to 200g/100L | | 14 days (H) | Apply at early bloom (1-10%), mid to full bloom (50-100%), petal fall and shuck fall, then at intervals of 14 days. Continue with a protective spray programme at 2 week intervals. * May be phytotoxic to Wilson, Wickson and Bellarosa varieties of plums. It is advisable to test any new plum varieties on a small scale before applying to the whole crop. |
| NON TREE AND VINE CROPS | | | | | |
| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
| Fruit Bananas | Leaf Spot | 2.2 to 4.5kg/ha plus spray oil (used according to its label directions) | 200g/100L plus spray oil (used according to its label directions) | NIL (H) | Apply when weather conditions favour disease outbreak. Repeat treatments In accordance with locally recommended spray schedules. Use sufficient water to ensure good coverage. NSW: <u>Ground application:-</u> Apply at 3 week intervals from December or January until May. <u>Aerial application:-</u> Apply at 7 to 10 day intervals from December or January until May. South QLD: Apply at 3 week intervals from December or January until May. North QLD: Apply every 2 weeks during the wet season. Extend to 3 weeks after this and then every 4 to 5 weeks during the drier months returning to 3 weeks prior to the next wet season. |

NON TREE AND VINE CROPS – continued

| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
|--|--|------------------|---------------------|--|---|
| Fruit Bananas – <i>continued</i> | Leaf Speckle, Cordana Leaf Spot, Fruit Speckle, Black Pit | 2.2kg/ha | 200g/100L | NIL (H) | Apply when weather conditions favour disease outbreak. Repeat treatments in accordance with locally recommended spray schedules. Use sufficient water to ensure good coverage. NSW: Ground application:- Apply at 3 week intervals from December or January until May. <u>Aerial application:-</u> Apply at 7 to 10 day intervals from December or January until May. South QLD: Apply at 3 week intervals from December or January until May. North QLD: Apply every 2 weeks during the wet season. Extend to 3 weeks after this and then every 4 to 5 weeks during the drier months returning to 3 weeks prior to the next wet season. |
| Vegetables Peas and beans grown for fresh consumption | | | | | |
| Broadbeans/ Faba Beans | Rust, Cercospora Leaf Spot | 1.7-2.2kg/ ha | – | 7 days (H) 7 days (G) | Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development. |
| | Chocolate Spot, Ascochyta Blight (suppression only) | 1.7-2.5kg/ ha | – | | If disease pressure is severe and weather conditions favour spread of the disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use the higher rate for dense crops and if disease pressure is severe. Volume of application: For ground apply 100L/ha; for aircraft apply at least 30L/ha. |
| Peas (<i>Pisum sativum</i>) | Rust | 1.7-2.2kg/ ha | – | | Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development. |
| | Ascochyta Leaf Blight (suppression only) | 1.7-2.5kg/ ha | – | | If disease pressure is severe and weather conditions favour spread of the disease, repeat the spray treatment at 2-3 week intervals in order to protect new growth and developing pods. Use the higher rate for dense crops and if disease pressure is severe. Volume of application: For ground apply 100L/ha; for aircraft apply at least 30L/ha. |
| Green Beans (<i>Phaseolus</i> spp.) | Angular Leaf Spot, Anthracnose, Rust | 1.7-2.2kg/ ha | – | | Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. Spray when leaves, flowers or pods show first signs of infection and weather conditions are likely to remain humid or wet and favour disease development. |
| Beetroot, Silver Beet, Spinach | Cercospora Leaf Spot, Downy Mildew | 1.7-2.2kg/ ha | 150 to 200g/100L | 14 days (H) | Spray at first sign of disease, then at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. |
| Capsicum | Target Spot | 2 to 3kg/ha | | 14 days (H) | |
| Carrots | Cercospora Leaf Spot, Alternaria Leaf Spot | 1.7-2.2kg/ ha | | 7 days (H) | |
| Celery | Septoria Leaf Spot | | | | |
| Cole crops: Broccoli, Brussels Sprouts, Cabbage, Cauliflower Cucurbits: Cantaloupe, Cucumber, Melons, Pumpkin, Squash, Zucchini | Alternaria Spot, Anthracnose, Downy Mildew, Ring Spot Downy Mildew, Anthracnose, Gummy Stem Blight, Septoria Spot | | | | |

| NON TREE AND VINE CROPS – continued | | | | | |
|-------------------------------------|--|------------------|--------------------|--|--|
| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
| Fennel (TAS only) | Leaf Blight (<i>Cercosporidium</i>) | 2.2kg/ha | – | 14 days (H) | Apply as a ground spray If disease symptoms are present in November, allow for one repeat treatment 10-14 days later. Aerial application may be required in prolonged wet periods. |
| Lettuce | Downy Mildew, Anthracnose, Septoria Leaf Spot | 1.7-2.2kg/ha | 150 to 200g/100L | | Spray at first sign of disease, then at 7-10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. |
| Onions, Garlic | Downy Mildew, Purple Blotch | 2.2 to 3.5kg/ha | 150 to 200g/100L | 7 days (H) | Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use higher rate and shorter spray interval during weather conditions favourable to disease development. |
| Potatoes | Early and Late Blight | 1.7-2.2kg/ha | 150 to 200g/100L | Not required | Apply when disease symptoms first appear and then repeat at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. |
| Rhubarb | Downy Mildew, Rust | | | 14 days (H) | |
| Tomatoes | Early Blight (Target Spot), Late Blight (Irish Blight), Leaf Mould, Anthracnose, Phoma Rot, Grey Leaf Spot | 2 to 3kg/ha | | 7 days (H) | |
| ORNAMENTALS | | | | | |
| CROPS | DISEASE/PEST | RATE | | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
| Carnations | Rust, Alternaria Leaf Spot | 150 to 200g/100L | | – | Apply when disease symptoms first appear and then repeat at 7-10 day intervals. Use higher rate and shorter intervals during weather conditions are favourable to disease development. |
| Chrysanthemum, Gladioli | Botrytis, Septoria Leaf Spot | | | | Caution: This product may be phytotoxic to some ornamental species, in particular some varieties of ferns. It is recommended to test a few plants before widespread use. |
| Roses | Black Spot | | | | |
| FIELD CROPS | | | | | |
| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
| Cotton Pima var only | Alternaria Leaf Spot | 2.5kg/ha | – | 14 days (H) | TITAN Mancozeb 750 WG Fungicide is a protectant fungicide. To maximize control ensure thorough coverage of all plant surfaces. Begin applications as soon as disease symptoms appear. Repeat before each infection period (eg dew or rain) but no later than 7 to 10 days after the last application. DO NOT apply more than 4 sprays per season. |
| Peanuts | Cercospora Leaf Spot, Peanut Rust | 1.7 to 2.2kg/ha | – | 14 days (H), 14 days (G) | Apply at interval of 10-14 days, starting when the disease symptoms first appear. If rust is anticipated apply at 7 to 10 day intervals. Use the higher rate and shorter spray interval during weather conditions favourable to disease development. |
| Poppies | Downy Mildew | 2.5kg/ha | – | 7 weeks (H) | Apply as a protectant spray when weather conditions favour disease development. Repeat spray at 14 day intervals if conditions persist. |
| PULSES GROWN FOR DRY BEANS | | | | | |
| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
| Broadbeans/ Faba Beans | Ascochyta blight, Chocolate Spot, Rust, Cercospora | 1 to 2.2kg/ha | – | 4 weeks (H) 14 days (G) | Spray when leaves, flowers or pods on plants first show signs of infection and weather conditions are likely to remain humid or wet and favour disease development. |
| Chickpeas | Ascochyta Blight, Botrytis Grey Mould | | | | If disease pressure is severe and conditions favour the spread of disease, repeat sprays at 2-3 weeks to protect new growth and developing pods. Use the higher rate for dense crops and if disease is severe. Repeat more frequently if severe disease persists. For improved coverage add a registered surfactant at 10-20mL/100L of spray. |
| Field Peas | Ascochyta Blight, Black Spot, Botrytis Grey Mould, Rust | | | | |
| Lentil and vetch | Ascochyta Blight, Botrytis Grey Mould, Rust | | | | |
| Lupins | Botrytis Grey Mould, Anthracnose | | | | |
| Beans (<i>Phaseolus</i> spp.) | Ascochyta Blight, Anthracnose Rust, Angular Leaf Spot | | | | |

PULSES GROWN FOR DRY BEANS – continued

| CROPS | DISEASE/PEST | RATE (per ha) | RATE (per 100L) | WHP DAYS Harvest (H) Grazing (G) | CRITICAL COMMENTS |
|-----------------|--|----------------------------|--------------------|--|---|
| Tobacco (Field) | Blue Mould, Brown Spot, Frog Eye Leaf Spot | 2.2kg/ha | 200g/100L | Not required | Field treatment: Apply at 7 day intervals commencing 10 days after transplanting and ceasing 7 to 10 days before the first harvest. Increase spray volume accordingly as plant size increases. Seed bed treatments: Apply when seedlings are 1.5cm across and then repeat twice weekly. Wet the seedlings evenly without run-off. |
| Turf | Helminthosporium, Fusarium, Rhizoctonia | 200-250g/100m ² | – | – | Apply as a protective spray schedule at 7 day intervals throughout the susceptible season. Use sufficient water to ensure adequate coverage. DO NOT feed grass clippings from treated areas to poultry or animals. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

Harvest

BANANAS: NOT REQUIRED WHEN USED AS DIRECTED.

MANGOES, PASSIONFRUIT: DO NOT APPLY LATER THAN 1 DAY BEFORE HARVEST.

CUSTARD APPLES AND PAWPAWS (PAPAYA): DO NOT APPLY LATER THAN 1 DAY BEFORE HARVEST.

PAPAYA LEAVES MUST NOT BE MADE AVAILABLE FOR HUMAN CONSUMPTION.

BEANS, CARROTS, CELERY, COLE CROPS, CUCURBITS, GARLIC, ONIONS, PEAS, SOYBEANS, TOMATOES: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

ALMONDS, BEETROOT, COTTON, CAPSICUM, FENNEL, LETTUCE, PEANUTS, POME FRUIT, RHUBARB, SILVERBEET, SPINACH and STONE FRUIT: DO NOT APPLY LATER THAN 14 DAYS BEFORE HARVEST.

GRAPEVINES: DO NOT APPLY LATER THAN 30 DAYS BEFORE HARVEST.

POPPIES: DO NOT APPLY LATER THAN 7 WEEKS BEFORE HARVEST.

GREEN LEGUMES, PULSE CROPS: GREEN CROPS FOR FRESH CONSUMPTION: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.

GRAIN CROPS FOR DRY BEANS: DO NOT APPLY LATER THAN 4 WEEKS BEFORE HARVEST.

CITRUS, POTATOES, TOBACCO: WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

Grazing

COTTON: DO NOT ALLOW LIVESTOCK TO GRAZE COTTON CROP, STUBBLE OR GIN TRASH THAT HAS BEEN TREATED WITH TITAN MANCOZEB 750 WG FUNGICIDE.

GREEN LEGUMES, PEANUTS, PULSE CROPS (grown for dry beans): DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER THE LAST APPLICATION.

FIELD PEAS AND BEANS (for fresh consumption): DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 7 DAYS AFTER THE LAST APPLICATION.

PAPAYA: DO NOT USE OR SUPPLY TREATED PAPAYA INCLUDING LEAVES FOR STOCK FOOD.

SOYBEANS: DO NOT GRAZE OR CUT FOR STOCKFOOD FOR 14 DAYS AFTER THE LAST APPLICATION.

GENERAL INSTRUCTIONS

TITAN Mancozeb 750 WG Fungicide is a protectant fungicide for the control of certain fungus diseases in many crops. For best results, apply as a regular spray programme.

Thorough coverage of the treated crop is essential.

APPLICATION

Dilute spraying: Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to point of run-off. Avoid excessive run-off. The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice. Add the amount of product specified in the Directions for Use table for each 100L of water. Spray to the point of run-off. The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

Concentrate spraying: Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed. Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume. Determine an appropriate dilute spray volume (see dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate. The mixing rate for concentrate spraying can then be calculated in the following way:

EXAMPLE ONLY:

1. Dilute spray volume as determined above: For example 1500L/ha
2. Your chosen concentrate spray volume: For example 500L/ha
3. The concentration factor in this example is: $3 \times$ (ie $1500L \div 500L = 3$)
4. If the dilute label rate is 10mL/100L then the concentrate rate becomes 3×10 , that is 30mL/100L of concentrate spray.

- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For further information on concentrate spraying, users are advised to consult the relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

For fruit trees and some row crops apply as a high volume dilute spray. The spray volume will vary according to the type of crop to be treated and should be increased in heavy growth and on large plants to ensure thorough coverage. The spray volume should also be increased as the size of treated crops increases throughout the growing season. As a guide for mature crops:

| Crops | Spray volume |
|---|------------------|
| Bananas, Tobacco, Vines and other row crops | 200 to 1100L/ha |
| Pome and Stone fruit | 1100 to 4000L/ha |
| Citrus | 4000 to 8000L/ha |

Aerial Application: May be applied by agricultural aircraft. Use at least 30 to 50L of spray mixture per hectare. (Refer Directions for Use table).

SPRAY TIMING

Treatments should begin prior to disease infection and continue until the threat of disease passes. Repeated applications may be required to protect new growth. Reduce the spray interval when weather conditions favour disease development.

MIXING

Slowly add the required amount of this product to the spray tank as it is being filled or thoroughly premix in a nurse tank for concentrate and aerial spraying. Add other products after TITAN Mancozeb 750 WG Fungicide is in suspension. Continue agitation while spraying to prevent product settling out and to ensure a uniform spray mixture. Rinse out spray tank, pumps, nozzles at the end of the day. When preparing spray solutions for use in a hand sprayer, premix as a slurry in a small container, and then add to sprayer containing one-third to half the desired final water volume.



COMPATIBILITY

This product may be harmful to *Typhlodromus pyri* and its use in orchards where integrated pest control is practiced should be avoided. TITAN Mancozeb 750 WG Fungicide is compatible with most commonly used insecticides and miticides such as Azinphos, Carbaryl, Dicofof, Endosulfan, Pirimicarb, Propargite, Sulphur Powder, Wetting Agent or Spreader-Sticker. Also compatible with DC-Tron. The user is advised to check the label recommendation for the companion product before preparing tank mixes.

Surfactants

The addition of a surfactant will improve initial spray deposits, fungicide redistribution and weatherability. Refer to manufacturers label instructions for further details.

FUNGICIDE RESISTANCE WARNING

Multi-site activity group of fungicides. For fungicide resistance management TITAN

GROUP **M3** FUNGICIDE

Mancozeb 750 WG Fungicide is a Group M3 Fungicide. Some naturally occurring individual fungi resistant to TITAN Mancozeb 750 WG Fungicide and other Group M3 Fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by TITAN Mancozeb 750 WG Fungicide and other Group M3 Fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, TITAN AG Pty Ltd accepts no liability for any losses that may result from the failure of TITAN Mancozeb 750 WG Fungicide to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT graze treated turf/grass; or feed turf/grass clippings from any treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to fish. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighbouring areas. DO NOT contaminate streams, rivers or waterways with the chemical or used container. DO NOT re-use container.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Shake and empty contents into spray tank/water/dip/drench, etc. DO NOT dispose of undiluted chemicals on-site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. DO NOT inhale dust. When opening the container and preparing spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves, goggles, and disposable dust mask covering mouth and nose. When using the prepared spray, wear cotton overalls over normal clothing, buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. Avoid giving alcohol.

SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for TITAN Mancozeb 750 WG Fungicide is available from TITAN AG Pty Ltd on request. Call Customer Service on (02) 9999 6655 or visit www.titanag.com.au

CONDITIONS OF SALE: TITAN AG Pty Ltd shall not be liable for any loss injury damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale supply use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on TITAN AG's skill or judgment in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of TITAN AG Pty Ltd has any authority to add to or alter these conditions.

EXPIRY DATE: 3 years after manufacture

Additional statements required by Globally Harmonised System of Classification and Labelling of Chemicals (GHS) and Safe Work Australia: May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. Very toxic to aquatic life. Precautionary: Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid breathing dust/fume/gas/mist/vapours/spray. Contaminated work clothing should not be allowed out of the workplace. Avoid release to the environment. Wear protective gloves/protective clothing/eye protection/face protection. IF ON SKIN: Wash with plenty of water. IF exposed or concerned: Get medical advice/attention. Specific treatment (see on this label). If skin irritation or rash occurs: Get medical advice/attention. Take off contaminated clothing and wash it before reuse. Collect spillage. Store locked up. Dispose of contents/container in accordance with local/regional/national regulations.

